

Variscan lithotectonic units in the Suchý massif of the Strážovské vrchy Mts., Western Carpathians – products of sedimentary, tectonometamorphic and granite forming processes


ĽUBOMÍR HRAŠKO, ZOLTÁN NÉMETH and PATRIK KONEČNÝ

State Geological Institute of Dionýz Štúr, Mlynská dolina 1, SK–817 04 Bratislava, Slovakia;
lubomir.hrasko@geology.sk

Abstract: The present geological investigation of the Pre-Alpine structure of the western part of Strážovské vrchy Mts (the Suchý massif; Western Carpathians) has distinguished three lithologically distinct Variscan lithotectonic units, which originated (1) in the deeper parts of oceanic basin (prevailing metapelites with different content of organic matter, metabasalts, metacarbonates); (2) sediments of the continental slope (flyschoid sediments with a predominance of greiwacke sediments; both VmD0); (3) a unit of continent basement primarily of pre-VmD0 granitic composition (orthogneiss). These rock sequences of differing geotectonic provenances were amalgamated and metamorphosed in the pre-intrusive (Pre-Mississippian) stage; pre-VmD2). Regarding the Variscan polyorogenic evolution, Variscan processes in Taticum represent the Meso-Variscan evolution (VmD).

The maximum P-T conditions of orogenic (regional) metamorphism (up to 610 °C and 7.5–8.5 kbar) were not sufficient for more extensive anatexis. Field observations indicated that the production of a limited volume of granitoid melts occurred mainly at the contacts of amalgamated lithotectonic units. Probably due to the heat, produced by the hot line below the collisional orogen and contributing also to unroofing processes (in orogenic phase VmD2) there started new granitization. This stage of Mississippian VmD2 granite formation is associated with the emplacement of various types of granitoid magma, encompassing the oldest granodiorites with frequent schliers, representing a poorly differentiated and poorly mobile crystal slurry (present in the SE part of the territory). In the highly evolved post-collision phase (post-VmD1 phase), masses of leucogranites intruded conformably with the deformation plan in the metamorphic complexes, interacting with surrounding metamorphic rocks in higher crustal conditions and causing their contact metamorphic overprint (up to 590 °C and 3–4 kbar). The syn-deformation character of leucocratic granites is proved by the orientation of biotite flakes parallel to the deformation plane of surrounding metamorphites. Part of leucogranites, especially in the central parts of the bodies, is omnidirectional. The stress field acted not only during the intrusion of the leucogranite magma, but also in the subsolidus stage. The final stage of this process is the formation of large bodies of pegmatites in extensional VmD2 fractures oriented at a large angle to the main stress component. The texture of the grey blocky K-feldspar pegmatites and the lack of H₂O-bearing minerals point to hydrothermal-pneumatolytic fracturing in a subsequently opened environment in VmD2. The final stage of VmD2 is represented by intrusions of I-type granodiorites with indications of magma mixing, or mingling. The chemical dating of monazites in granitoids allowed to date individual phases of granitization process in the range of 360–345 Ma, which youngest ages corresponded with the formation of pegmatites. Dating of monazite in metamorphic rocks points to their thermal overprint during granitization process (360–350 Ma), having already earlier metamorphic overprint (380–370 Ma). The scenario of placement of granitoid intrusions is consistent with the decompression regime (in VmD2) after the end of VmD1c crustal thickening until the fracturing of the crustal block with of I-type magma intrusions of deeper origin. After this period, the exhumation of crystalline blocks, partial diaphoresis and later surface erosion continue until the Lower Triassic. Re-submergence of the crystalline complexes is associated with a low degree of Alpine metamorphic reworking.

Graphical abstract



Highlights

- Three types of Meso-Variscan sequences (lithotectonic units; LTUs) were distinguished in the Tatic crystalline basement of the western part of Strážovské vrchy Mts (the Suchý massif), documenting its complicated Paleozoic geological evolution: (1) LTUs of the Lower Paleozoic riftogenic oceanic sedimentation, (2) flyschoid sediments (Paleozoic continental slope sediments; both VmD0) and (3) their basement crustal rocks – orthogneisses (originally pre-VmD0 granitoids). These complexes were amalgamated in a collisional event VmD1c and later at granite forming processes VmD2. The VmD2 metamorphism has produced an initial melting. The formation of granitoids in decompression phase produced thermal overprint of the amalgamated complexes at the granite front. In the first stage, less mobile crystal slurries were formed, which inherit the structural characteristics of the surrounding metamorphites. In the second phase, the complexes were sheared and two-mica granites and pegmatites were formed; this additional thermal metamorphism overprinted also earlier metamorphosed complexes. In the last stage of VmD2 evolution, I-type granodiorites, tonalites and diorites indicating magmatic mixing have intruded. CHIME monazite dating indicates the age of granite formation of 360–345 Ma.

Introduction

In 2015–2020 the State Geological Institute of Dionýz Štúr in Bratislava (Slovakia) carried out new geological mapping and regional geological research in eastern part of the Strážovské vrchy Mts, resulting in compilation of a new regional *Geological map of the Strážovské vrchy Mts (eastern part) at a scale of 1 : 50 000* in 2021 and explanatory notes to this map (currently edited). New map represents an update of a part of earlier geological map compiled by Maheľ et al. (1981). New research focused on Alpine tectonic units: Tatricum, Fatricum, Hronicum and younger geological formations (Paleogene, Neogene and Quaternary). This article is devoted to Variscan evolution

and geological structure of the Tatricum crystalline basement of the western segment of mapped region, geographically belonging to the Suchý massif. Despite defining Meso-Variscan lithotectonic units (LTUs) in the Suchý massif, we briefly classify in studied area also those of Cadomian, Paleo-Alpine and Neo-Alpine orogenic cycles.

General characteristics of crystalline basement and its previous researches in the Strážovské vrchy Mts

First and the most comprehensive research of the Strážovské vrchy Mts crystalline basement by A. Klinec, I. Lehotský and M. Ivanov (Klinec, 1956, 1958; Ivanov,




Fig. 1. Tectonic scheme of the Strážovské vrchy Mts, with tectonic blocks of Tatricum (crystalline basement and tectonically reduced Mesozoic sequences of the Malá Magura Unit), nappes of Fatricum and Hronicum, as well as Cenozoic sediments. Variscan crystalline basement (VmD) is divided by N-S trending Neo-Alpine (AnD4) Diviaky fault (D) to Suchý and Magura massifs. The Suchý massif is tectonically rimmed by AnD34 Zdychava fault (Z) and Brekovský potok fault (B) and further segmented by intracrystalline Radiša fault system (R) with more pronounced Neo-Alpine deformation of crystalline formations.

1957) included mapping of this region in the 1950s for the sake of compiling its first general geological map at a scale 1 : 200 000.

Ivanov (1957) parallelized the crystalline basement of Suchý and Malá Magura massifs with SW continuation of the Malá Fatra crystalline basement, formed of inhomogeneous sequence of crystalline schists and granitoids. Author correctly recognized the protoliths of metamorphites, as well as multi-stage magmatic granite genesis. The genesis of granitoid rocks he affiliated with Variscan (earlier used synonym used by some authors Hercynian) late- to post-orogenic magmatic cycle. He considered the migmatites, present in paragneisses, mainly as thermodynamic product of granitoid intrusions.

Detail mapping and petrographic research of crystalline basement in the 1970s by Kahan et al. (1978) and Kahan (1979) distinguished several types of metamorphic and granitoid rocks, encompassing also their structural research.

Petrography and geochemistry of amphibolites and associated black schists (graphite gneisses) in Malá Magura and Suchý massifs were investigated by Ivan and Méres (2015). They equate them with effusive basalts, dolerites and gabbros as well as deep water sediments, rich in silica and carbon, being hydrothermally altered before orogenic metamorphism. These rocks represent upper part of ofiolite suite, equal with that, present in Upper Devonian Pernek Group of the Malé Karpaty Mts. The finding of distinctive Cr, V and Mn rich metamorphic rocks and Ca garnets in black schists in the Čierna Lehota area supports this opinion (Bačík et al., 2018). In the area of the Závada valley north of the Závada pod Čiernym vrchom village, the exploration activities were carried out in black schists for graphite raw material (Mikoláš et al., 1995).

Hovorka and Méres (1991) geochemically investigated paragneisses in both crystalline cores of the Strážovské vrchy Mts and also in the Malá Fatra Mts. They interpreted low K/Na lithologies of the continental crust as their source material (especially intermediate to acid magmatites of granodiorite-tonalite composition). The sedimentary maturity of the original material is the highest in the Suchý massif and is manifested by a higher content of the clay component.

Putiš (1976) and Kahan et al. (1978) comprehensively evaluated the structural elements of the crystalline basement (Malá Magura and Suchý massifs) of the Strážovské vrchy Mts, including microstructural analysis. They distinguished secondary foliation planes s1 of metamorphic origin (crystallization schistosity), present in paragneisses and simultaneously forming biotite schliers in granitoids. The younger s2 planes, bearing retrograde facies minerals (sericite, chlorite, bauerite, albite, quartz), penetrated s1 planes. Planes s3 of retrograde metamorphism (according to Kahan et al., l.c.) were formed at arching

of crystalline cores near the contact with Mesozoic and Cenozoic complexes. Work evaluates also fold structures in crystalline basement, mineral lineations and joints, identifying directional block displacements.

Based on microstructural research of preferred quartz orientation, P. Kováč (1986) demonstrated that in the Suchý massif granitoids quartz elongations are trending NNE–SSW, correspondingly with the course of granitoids, as well as foliation in metamorphites, described by Kahan (1978). P. Kováč (l.c.) supposed that preferred orientation of granitoids, investigated in quartz and micas, represent a one-act process, manifested by recrystallization and rotation of quartz and mica minerals.

Based on drilling works and surface exploration trenches in the Čavoj area (Mikoláš et al., 1995), the synform and antiform geological setting of paragneisses was revealed with relatively steeply dipping fold planes and intensive mylonitization of the crystalline basement in the wells increased N-ward, i.e. towards the contact of crystalline complexes with the Mesozoic cover (Malá Magura Unit), formed by quartzite at their contact. Stronger Alpine overprint along the margin of crystalline massif was proved by this way.

Based on relationship of metamorphic rocks, their metamorphic grade, granitization and migmatization, Kahan (1979, 1980) states two alternatives of thermal overprint – either by anatexis process, i.e. orogenic metamorphism producing anatexis, or overheating of “supracrustal” metasedimentary series.

Investigating metamorphic zonality in Suchý massif, Korikovski et al. (1987) distinguished two metamorphic zones: staurolite-andalusite-biotite at the margin and sillimanite-biotite-muscovite in remaining part of crystalline basement. The petrogenetic grid revealed PT conditions of metamorphism 3 kbar (300 MPa) and 540–590 °C. Muscovite-sillimanite aggregates in peraluminous granites indicate their origin at the expense of magmatic feldspars and biotites through the loss of Mg, Fe, Na, partly K, which, according to authors, points to the process of high-temperature acid leaching. Simultaneously they separated this process from pervasive muscovitization in connection with granites. Authors (l.c.) consider this process to be a deeper analogue of greisenization with metasomatic origin of muscovite and andalusite.

The presence of the staurolite-sillimanite and garnet mineral association especially at the W margin of the Suchý massif Dyda (1988) explains by the pre-metamorphic protolith rich in clay schists. According to Dyda (1990), mineral balances indicate different metamorphic conditions in the Malá Magura and Suchý crystalline basements: In the first case their culmination reached 640 °C with pressure of 5 kbar and in the case of the Suchý paragneisses it was 560 °C and pressure of 4.5 kbar.

Hovorka and Méres (1991) distinguished two events in metamorphic overprint of the paragneisses of the Strážovské vrchy and Mala Fatra crystalline basements. The first event – found only in the Malá Fatra basement (garnets with pyrope content above 30 %) – took place in higher temperature amphibolite facies (sillimanite zone), second event – under lower conditions (staurolite and sillimanite zone) – was revealed in both basements.

High-temperature metamorphic conditions were determined by TERMOCALC software (Čík & Petřík, 2014) from migmatite and paragneiss of the Magura massif (Poruba valley). The peak conditions for migmatite – 782 ± 53 °C (resp. 670–760 °C), pressure 7.4 ± 1.7 kbar (resp. 6.9–7.4 kbar). For paragneiss it was 668 ± 53 °C (resp. 700–770 °C) at pressure 5.5 ± 1.2 kbar (resp. 6.7–8.2 kbar). Younger low-temperature metamorphism was defined in the Magura massif (l.c.) by the presence of margarite and pumpellyite. Calculations of retrograde conditions in migmatite gave values of 480 °C and pressure of 4.6 kbar and in paragneiss 300 °C and a pressure of 2.9 kbar.

The composition and ages of granitoids and paragneisses from the Strážovské vrchy Mts. were studied by Viliňová (1988, 1990). She divided tonalities, granodiorites and granites based on their petrochemical and mineral characteristics. Rocks of granodiorite composition dominate in the Suchý massif, less often are tonalities and granites. In the gneiss complex she (l.c.) investigated the fine-grained gneisses, banded gneisses and augen gneisses [in present geological map by Hraško & Kováčik (eds.) et al., 2021, being redefined to orthogneisses]. Viliňová (l.c.) investigated also chemical composition of the trace elements in feldspars and stated their structural arrangement. Fe²⁺ biotites in all granitoid types with a relatively small compositional range, correspond by composition with biotites from gneisses. In garnets of granitoids, which are relatively non-zonal, the content of the spessartite molecule increases from leucogranites to granodiorites and tonalites. The normalized REE curves show the absence of Eu minimum in tonalites and enrichment of light REEs. The application of feldspar Plg-Kfs thermometry points to subsolidus equilibrium temperatures in granitoids (up to 550 °C). Leucogranites of the Strážovské vrchy Mts are frequently bearing garnets of several mm up to a few cm in size, which occur together with sillimanite, biotite, muscovite in the prevailing quartz–K-feldspar–plagioclase granite composition. Hovorka and Fejdi (1983) distinguished two types of garnets with very close composition. Garnets have prevailing almandine component (1. Alm₇₁, Py_{9,5}, Spess₁₈, Gross₁, Andr_{0,5} and 2. Alm₇₅, Py₁₂, Spess₁₀, Gross_{0,5}, Andr_{2,5}). Authors interpret this difference in composition to be the result of melt crystallization under different thermodynamic conditions – they crystallized before the emplacement of magma into higher crustal conditions.

Král et al. (1987) investigated isotopic composition and age of granitoids based on ⁸⁷Rb/⁸⁶Sr isotopic system, including more types of granitoid samples and aiming to calculate the isochrone Rb/Sr age from the Suchý and Malá Magura massifs. Isochrone age was calculated to 393 ± 6 Ma, with initial ⁸⁷Sr/⁸⁶Sr ratio 0.7060 ± 0.0002 . According our opinion these samples were not genetically related. The dating of S-type granite sample (V-94) by Král et al. (1997) using TIMS analysis provided the age of 356 ± 9 Ma.

More recent zircon dating based on SHRIMP analysis (concordia age) from sample of S-type muscovite-biotite granite with garnet (sample MM-29) performed by Kohút & Larionov (2021) from the Magura part of the Strážovské vrchy Mts and from the Porubský potok stream provided the age 360.9 ± 2.7 Ma.

The geological structure of the territory was comprehensively described in the monographs of Mahel' (1983, 1985).

Used methodology

The field geological research was supported by detail petrography of mapped rock types. Chemical composition and BSEI images of rock-forming minerals were obtained by electron microprobe CAMECA SX 100, housed in the State Geological Institute of Dionýz Štúr (P. Konečný, V. Kollárová). In the laboratory, monazite dating methodology MARC (monazite age reference calibration) developed over past years by P. Konečný (2018 including further improvements) was applied to granitic rocks.

Monazites suitable for dating have to meet several conditions. A low concentration of Pb in the order of tenths of ppm can only be detected with the application of a strong analytical conditions (e.g long counting times and high beam current). Pb was acquired with accelerating voltage of 15 kV, long counting times of 300 s for peak and 150 s for two background points were applied and the high sample current of 180 nA was used. The electron beam diameter was adjusted to 3 µm. Under such conditions, the PbMα line is effectively excited and offers acceptable compromise between beam damage and counting efficiency. Th, U included in the age calculation and Y overlapping the PbMα were also precisely measured with an extended measurement time of 35 s, 90 s and 45 s, respectively. The following calibration standards (natural minerals or synthetic components) were used to calibrate the electron microprobe: apatite (P-Kα), wollastonite (Si-Kα, Ca-Kα), GaAs (As-Lα), baryte (S-Kα, Ba-Lα), Al₂O₃ (Al-Kα), ThO₂ (Th-Mα), UO₂ (U-Mβ), cerussite (Pb-Mα), YPO₄ (Y-Lα), LaPO₄ (La-Lα), CePO₄ (Ce-Lα), PrPO₄ (Pr-Lβ), NdPO₄ (Nd-Lα), SmPO₄ (Sm-Lα), EuPO₄ (Eu-Lβ), GdPO₄ (Gd-Lα), TbPO₄ (Tb-Lα), DyPO₄ (Dy-Lβ), HoPO₄ (Ho-Lβ), ErPO₄ (Er-Lβ), TmPO₄ (Tm-Lα), YbPO₄ (Yb-Lα), LuPO₄ (Lu-Lβ), fayalite (Fe-Kα) and SrTiO₃ (Sr-Lα). UMβ overlapped by ThMζ, ThM3-N4, ThM5-P3 and PbMα overlapped by ThMζ1, ThMζ2 and PbMα overlapped

also by $Y_{L\gamma 2,3}$ as well as numerous other interferences among REE were corrected by correction coefficients obtained by measurement on calibration standards. Each analysis of monazite includes the complete set of elements of which it is composed. We are avoiding the methodology where only Th, U, Pb and Y are measured and the other elements are measured elsewhere in the same monazite zone. Such method creates inaccuracies depending on the homogeneity of the monazite and also on the matrix effect, where all elements are required for the ZAF correction. The age determined from the given analysis is only the apparent age. To find out the real age, it is necessary to combine measurements at several points (10-50). The age group plotted on the histogram can indicate whether it represents one or several age populations. The resulting age from the age of population is then calculated using the weighted average of the apparent ages. On the isochron (Pb vs. Th – including the contribution from U) the age population is spread around a linear relationship. The linear regression should ideally intersect the zero coordinate and thus represents an additional test of the correctness of the dating. The method of age calculating based on statistical evaluation was described by Montel et al. (1996). The methodology developed in the Electron Microanalysis Laboratory (ŠGÚDŠ) represents an improved methodology that is practiced only in this laboratory. It is based on a final correction using 7-9 monazite age standards, where the age is determined by more accurate isotopic dating methods – Th-U-Pb: SHRIMP, LA-ICPMS, ID-TIMS, etc. Methodology named as MARC (Monazite Age Reference Correction, P. Konečný et. al., 2018) includes corrections for interferences, corrections for exponential background, an innovative method of determining the dependence between the composition of monazite (average atomic number) and the difference between the linear and exponential background for the $Pb/M\alpha$ line, determining the dependence between the measured Pb and the theoretically required (ΔPb) to reach the age of the monazite standard, determination of MARC coefficients, which will be used for final fine-tuning of the age calculation and others.

The pressure–temperature conditions of metamorphic overprint were determined by R. Demko (SGUDS), using methodology of garnet-biotite-plagioclase-quartz (GBPQ) geobarometry in medium- to high-grade metapelites (Wu et al., 2004).

Classification of lithotectonic units – LTUs – is based on orogenic (Wilson) cycles, being indicated by the **XD labelling method** in polyorogenic terrains in the prefix X (cf. Németh, 2021; Fig. 2 *ibid*), as well as by affiliation of individual orogenic phases of these cycles by a **number after D**: **XD0** – divergent process of riftogenesis; **XD1** – convergent processes of subduction (XD1s), obduction (XD1o) and closure of elongated oceanic space by collision (XD1c); **XD2** – post-collisional thermal / deformation processes, unroofing and metamorphic core

complex evolution; **XD3** – intraplate consolidation – strike-slip faults preferably of NW–SE and NE–SW trends (cf. Németh et al., 2023), transpression, transtension, rotation of blocks, *etc.*; and **XD4** – regional extension (pure shear-type regional faults, dominantly of E–W and N–S courses, cf. *l.c.*). From **X prefixes** for orogenic cycles within Europe suggested by Németh (2021) in this paper we use: Cd – Cadomian, V – Variscan, Ap – Paleo-Alpine (Mesozoic Alpine orogenic cycle) and An – Neo-Alpine (Cenozoic Alpine orogenic cycle; both represent complete orogenic cycles).

Similar principle of designation is used at metamorphic overprints – the **MX labelling** (e.g. MV0, MV1sc, MAp2, etc., *l.c.*)

New characteristic of lithotectonic units of the pre-granite crystalline basement in the Suchý massif

Published geological map by Hraško & Kováčik (eds., 2021) for the reader of this paper is available on:

https://www.geology.sk/wp-content/uploads/documents/foto/geol_mapy_50k/58_StrazovskeVrchy_final.jpg.

Detail geological mapping of crystalline basement in the western part of Tatric occurrences in the Strážovské vrchy Mts (Suchý massif), research of structural and spatial relations acting before the granite-forming process and lithological peculiarities, allowed to define three below stated main lithotectonic units (LTUs) of differing geotectonic provenience. Rock sequences of these LTUs were amalgamated during Variscan collision and later partial melting.

1. **LTU of deep-water euxinic facies (Vmd01s)**, in pre-metamorphic state consisting mainly of fine-grained pelitic and quartz sedimentary facies rich in organic matter, with preserved fragments of accompanying oceanic mafic volcanites;
2. **LTU of proximal sedimentary facies of continental slope (Vmd01s)** of Paleozoic basin, built mainly of flyschoid sediments of sandy-greiwacke composition. In the field, these can be distinguished by alternating of metamorphosed sandy vs. clayey-sandy beds (less Al-rich sediments), having lower content of organic matter and rare occurrence of small metabasite bodies;
3. **LTU of orthogneisses** is common in the Suchý massif. In smaller scale orthogneisses occur also in the W part of the Magura massif. They represent a crustal granitoid segment thrust at Meso-Variscan collision (**Vmd1c**) on previously described **Vmd01s** LTUs. Later the originating thick skinned sequence was metamorphosed in **Vmd2 / MVm2**. This LTU of orthogneisses crops out in upper structural position of fan structure with fan axis

trending ENE–WSW (shown in geological profile 1–2 in published geological map by Hraško & Kováčik, eds., 2021, l.c.).

These originally separated complexes were interconnected and submerged owing to convergence in zone of elongated Variscan basin. In the area of the lower part of the metamorphic packet and along shear faults, a partial melting of the protolith with an appropriate lithological composition took place in **VmD2**.

General lithological and metamorphic characteristic of new lithotectonic units

Lithotectonic unit built of deep-water euxinic facies (VmD01s)

It is originally formed mainly of the fine-grained pelitic and quartzly sedimentary facies rich in organic matter, with preserved fragments of accompanying oceanic mafic volcanites;





Fig. 2. Association of amphibolites (123) and black schists (119) SE of the Čierna Lehota village. Numerous mining works exploring possible sulphide mineralization are typical for this area.

Fig. 3. Amphibolite bodies (123) and small bodies of tremolite-plagioclase amphibolites (124b), black graphitic schists, black paragneisses and black metaquartzites (119), lying on the complex of schistose or massive biotite and two-mica paragneisses (113) (continental slope sediments). Orthogneisses (111) occur in upper structural position. Geological situation near the Závada stream.






Fig. 4. The position of the largest amphibolite body (123), concordantly with the subvertical deformation structure, at the contact between migmatites with prevalence of stromatitic to pygmatitic textures (117a) and massive to weakly oriented hybridic (schliered) granodiorite with frequent biotite schliers and xenoliths of paragneisses (110). No black schists were found, but according to presence of remnants of old mining in the northern part of body, their presence is probable. Amphibolite has signs of partial melting and intrusion of later leucogranites. Situation between Krstenica a Nitrica valleys.

Amphibolites and associated rocks (amphibolic and amphibole-biotite gneisses, quartz massive paragneisses, graphitic metaquartzites, pyroxene amphibolites, amphibolites with garnet and tremolite-plagioclase amphibolites) are in earlier geological maps from this area (Kahan in Mahel' et al., 1982; Ivanov, 1957) shown in summary as amphibolites. Visualized are mainly large bodies at the Chvojnica area (Magura part), Liešťany (S part of the Krstenica valley) and in area north of the Závada pod Čiernym vrchom village (Závada valley, area of its right tributary – the Železná dolina valley). Amphibole-bearing rocks and associated sediments with prevailing pelitic / resp. quartz composition represent common source of disseminated sulphidic mineralization, which is demonstrated also by numerous old mining works in their space. Also based on the remnants of old mining we have identified such lithologies also in the westernmost termination of crystalline basement in the Suchý massif (SE of Čierna Lehota village, area of Závadská poľana hill).

Due to metamorphic overprint, it was not possible to express quite well the facies differences between the metapelitic and quartz-pelitic, siliciclastic metasediments in cartographic visualization. Despite, in wider context in areas with higher representation of amphibolitic rocks, there was observable the presence of metasediments with higher content of Al-rich minerals – sillimanite and staurolite, as a reflection of more pelitic composition of metasediments. In the surrounding of smaller mafic bodies in area located SW and NW of Suchý vrch hill (altitude point 1028), there occur also metapelitic metamorphites with common macroscopic staurolite, representing an indicator of Al-rich protolith. At the contact zone of stromatitic migmatites (117a) and massive to weakly oriented schlier granodiorites (110) in the area of Nitrica river and Krstenica valley there were found two smaller and one big bodies of metamafites (123), which underwent interaction with leucogranitic melt (Fig. 4).

Amphibole occasionally occurs also in paragneisses, which can be in this case named as “amphibole-biotite gneisses”. These thin lithological intercalations we do not describe here, but in geological map some their occurrences are visualized with a special hatch (122).

The amphibolite bodies occur in the Suchý massif concordantly with the structural elements of the surrounding metamorphites.

Present types of amphibolic rocks

Amphibolitic rocks with calcareous-silicate component, locally with garnet and clinopyroxene (124a)

They are usually heterogeneous, often complexly deformed rocks, appearing in various green, greenish-grey shades in the area of larger amphibolite bodies of Suchý hill.



Fig. 5. Dark-grey amphibolite rock with thin light-grey interbed of diopside pyroxene (area of Železná dolina valley, sample SZN-32a).




Fig. 6. Microphotographs of polymetamorphic clinopyroxene-amphibole amphibolite (sample SZN -32a, area of the **Železná dolina** valley, being a tributary to the Závada valley). A–B – granoblastic diopside (highlighted by red arrow) from Ca-rich, originally carbonate interbed in tuffitic sequence; C–D – dynamic deformation of older plagioclase (highlighted by black rectangle) with S-C setting represents younger deformation phase in the epidotic amphibolite facies.




Fig. 7. BSEI of clinopyroxene-amphibole amphibolite, sample SZN-32a, the **Železná dolina** area. A – Assemblage of diopside (cpx) with amphibole (tschermakite and younger actinolite), accessory apatite and chalcopyrite. B – Rotation of older plagioclase due to the decomposition of former plagioclase to Pl X_{An} 40–48 %, muscovite, zoisite and rim consisting of grey actinolite. Light phases represent titanite. Assemblage in part of this rock, consisting of prehnite, albite and actinolite belongs to the lowest thermal part of polymetamorphism.

Their affiliation to metamorphosed calcium-silicate rocks is evidenced by the increased proportion of epidote (clinozoisite), the common presence of clinopyroxene phenocrysts and garnet rich in grossularite component. The plagioclase-amphibole-dominated composition of the rocks and the loose spatial association with amphibolites indicate that the primary lithology consisted of pyroclastic (?) products of basic volcanism enriched with a calcareous admixture. Microscopically present bright positions parallel to the metamorphic foliation are formed by diopside (Figs. 5 and 6).

The composition of clinopyroxenes corresponds to diopside with a content of 1.3 % jadeite component, older amphiboles correspond to tschermakite, while younger recrystallized amphiboles are of actinolite composition.

Plagioclase is usually recrystallized to form a new generation $X_{An} = 41-48\%$. Based on the presence of clinozoisite, which was formed at the expense of plagioclase, the original plagioclases were more basic. The lowest thermal part includes the albite-prehnite association, which is probably of Alpine age.

Tremolite-plagioclase fine-grained pale amphibolites (124b)

Rarely present light fine-grained rocks are part of the complex of amphibolites (metabasalts) and appear at the contact with black metaquartzites, irregularly rimming the southeastern part of the elongated body of amphibolite in the area of the Železné and Závada valleys north of the village of Závada pod Čierny vrchom.


Considering the mineral composition, they probably represented mica-carbonate (dolomite) sedimentary inlays in black metaquartzites in contact with metabasalts. The alternative of altered ultramafite due to the lack of sedimentary textures is also possible (this would also be indicated by the position in the lower part of the amphibolite

body, which could represent a fragment of the ophiolite suite). Another alternative is that the rock represented the acidic part of the ophiolite suite (keratophyre).




Fig. 8. Sample of tremolitic-plagioclase rock with deficiency of dark minerals. In the field this rock resembles the fine-grained quartzitic rock (sample SZN-24).

The samples represent a light omnidirectional fine-grained rock (Figs. 8 and 9), without dark minerals. The microscopically visible metamorphic structure of the rock is formed by omnidirectional colorless strips of amphibole of tremolite composition. The basic material is composed of plagioclase (An_{41-43}). Younger metamorphic overprint is




A




B


Fig. 9. A, B – Microphotographs of sample SZN 24 from the area between the Závada and its right-side tributary from Železná dolina in western part of the Suchý massif. A rarely occurring amphibolic (tremolitic)-plagioclase fine-grained rock at first glance resembles metaquartzite. It contains a ground mass formed by plagioclase and phenocrysts of light amphibole of tremolite composition. The interstices between amphiboles of tremolite composition are formed by plagioclase. This rock occurs in marginal parts of large amphibolite bodies. A – parallel (N//) and B – crossed nicols (NX) of polarizing microscope.



A



B



C

Fig. 10. Microphoto of the sample LHS 261: A, B – ne-matoblastic structure of rock borned by tremolitic to Mg-hornblende amphibole and plagioclase, N//, NX. C – BSEI – besides tremolitic to Mg-hornblende amphibole and plagioclase (An – 32, 46, 61 %) rock contains a great number of sulphides – pyrite, pyrrhotite, chalcopyrite. Younger, lower temperature metamorphic association is formed by albite, chlorite and pumpellyite.

associated with formation of chlorite, barite, pumpellyite and oligoclase (An₂₈).

A similar sample LHS-261 from the Suchý massif taken from Závada and Železná dolina valleys area at the junction of paragneisses and black-graphitic schists in continuation of the amphibolite body above E side of the Závada valley. The rock resembling fine-grained light quartzite contains tremolitic-Mg-hornblende amphiboles, basic plagioclase of andesine-labradorite composition (max. An 61 %). The presence of sulphides is common.

Amphibolites – olive-green-grey fine-grained foliated types (123)

Amphibolite bodies were already considered by Ivanov (1957) to be part of the ophiolite (oceanic) suite of basic magmatites, which was confirmed by later geochemical research.

Amphibolites usually form several meters, max. the first 10 m thick bodies of dark green to grey fine-grained, detailed foliated rocks, mostly conformal with the surrounding paragneiss substrate. Tiny lenses of amphibolites are irregularly dispersed in the metamorphic substrate.

In the Suchý massif there are two to three more distinct bodies, or groups of bodies. The easternmost of them is located in the Liešťanská dolina valley (Krstenica valley), and like the body in the Chvojnica valley in the Magura part of the crystalline basement, it occurs at the junction of migmatites and granitoids, with exceptionally preserved higher-temperature paragneisses in the vicinity. From the point of view of the Variscan setting, this body occurs in the lower tectonic position at the contact of the lower part of the schist-like granodiorites near their contact with the migmatites (Fig. 4). These bodies are probably the relics of a larger set of metapelites, metapsammites, which, however, are found only rudimentarily in this position in the form of paragneisses. Only rarely is it possible, e.g. E from Nitrica to find amphibolite relics in association with paragneisses in hybrid granitoids. Such a lack of metasedimentary associating lithologies is very likely a consequence of the tectonic reduction of a significant part of the volcanic-sedimentary complex during the granitizing Devonian-Lower Carboniferous event. The second position of the amphibolites in the Suchý massif is the upper structural position of the amphibolites, which emerges both in the area of the Závada valley and its tributary Železná dolina valley, and on the other hand from the ridge of Závadská poľana towards the Čierna Lehota village, where the amphibolites are part of a complex of black schists (graphitic metaquartzites), graphitic and graphite-amphibole paragneisses. The association of amphibolic rocks with metapelitic rocks,

which are characterized by Al-rich metamorphic minerals – sillimanite and staurolite, is striking in the western part of the territory. This structural position is spatially more remote from granitization processes.




Fig. 11. Banded amphibolite from the zone of granitization (sample LHS-153) above left side of the Krstenica valley (Liešťanská dolina valley) under this high voltage line. Rock in brittle regime is intruded by aplitic leucogranite without mutual interactions, which proves intrusion of leucogranite in higher crustal level with more rapid crystallization of leucogranite magma.


The mineral content of amphibolites is usually characterized by a significant predominance of common amphibole with green pleochroism prevailing over plagioclase. In minor quantities, but characteristically represented minerals are titanite, epidote, or clinozoisite and opaque ore phases. As a rule, amphibolites are fine-grained (the average grain size reaches around 0.2 mm), amphibole in some places makes up to 90 vol. % of the mineral composition of the rock.

Amphibole-biotite and amphibole-plagioclase gneisses (122)


Due to the problematic differentiation of amphibolic gneisses and amphibolites at field mapping, these are usually cartographically included under the group of amphibolites (123). Only in some places do we highlight in the geological map with a special symbol the areas with a greater occurrence of paragneisses with amphibole, in which biotite and amphibole gneisses alternate, which often contain several cm thick positions richer in biotite and positions richer in amphibole. The marginal member is represented by amphibole-plagioclase gneisses, which originally probably represented intercalations of basic tuffites in the marine sediment. Plagioclases (from the




A



B



C



D


Fig. 12. Transitional type of paragneiss. Qtz-Plg-Hbl-Bt-Grt-Chl paragneiss in the vicinity of amphibolite bodies (sample SZN-57) contains often graphitic pigment. A – N//, B – NX: microphoto of lepidogranoblastic structure of paragneiss with often graphitic pigment. C, D – BSEI. D – Hornblende (Hbl) closes irregular garnet (Grt).

sample SZN-85 from the Závada valley area) correspond in composition to the oligoclase / andesine interface ($X_{An} = 31\%$). Amphibole corresponds to chermakite to magnesiohornblende, and the younger generation corresponds to actinolite.


Graphitic schists, graphitic paragneisses and black graphitic metaquartzites in association with amphibolites (119)

Dark grey schists with an organic, graphite substance are sporadically scattered within the metamorphic substrate, conformly with its metamorphic schistosity. Distinctive beds of black schists have been identified near several amphibolite bodies, suggesting a genetic link between basic volcanism and euxinic facies sedimentation in deep-water conditions. Such rocks are also present in the SE region from the elevation Klin (769 m) towards the Jasenina stream, or, according to the drilling works, they are also part of the paragneiss complex, where amphibolites occur only rarely (area SW from the settlement of Gápel towards the settlement of Petriská). The spread of graphite-rich lithologies in the paragneiss complex is relatively frequent, but without detailed surface petrographic or geophysical


research, it is difficult to visualize it spatially. Graphitic rocks are finer-grained, contain more quartz and muscovite, which are otherwise insignificantly represented in the metamorphic substrate in the primary metamorphic form. These rocks are generally poorer in plagioclase and biotite and primarily represented black schists. They can also be considered to be suitable carriers of disseminated sulphide mineralization, as indicated by remnants of old exploration activity in graphitic or silicified rock environments. At the scale of the geological map, identifiable occurrences of these lithological variations, similar to the gneisses with amphibole mentioned below, are marked with a special hatch. Dark grey biotite paragneisses are also an example (sample LHS-350 – Fig. 13) with an organic admixture in association with black graphitic schists and amphibolic schists in the area of the westernmost termination of the crystallinics above the right side of Brekovský potok valley (south from the Čierna Lehota village). Paragneisses have a lepidogranoblastic structure with a predominance of Qtz-Pl matter, Bt has metamorphic preferred orientation, associated in places with zonal Grt, which has cloudy edges (the poikilitic edge part is a consequence of new thermal metamorphism). Very fine organic matter is



A




B




C

Fig. 13. A, B – Microphotograph at various magnifications (massif N, SSE from the Čierna Lehota village, sample LHS-350): A – view on structure of Bt paragneiss with fine disseminated graphite bound mainly on plagioclase, N//; B – detail of association Qtz-Plg-Bt-Grt; C – BSEI of garnet in assemblage with Bt and Chl in plagioclase. Garnet closing poikilitically fine Qtz-Pl-Chl, inclusions are richer present in marginal part of garnet, which causes the cloudy character of the Grt edges.



A



B

Fig. 14. A, B – Sample SZN-25 from the area of Železná dolina valley; A – greyblack fine-grained cevernous metaquartzite rich in graphite, porous after the weathering of sulphidic impregnation; B – N//, **C–NX** Thin section of the rocks composed of from a substantial proportion of organic matter and quartz.




Fig. 15. Position of migmatitized paragneisses (116) with small content of ductile folded leucosomes, neighbouring with migmatites (117a) and schlier granodiorites (110).



Fig. 16. Migmatitized paragneiss with distinct primary “bedding of metasediment” $s_0 = s_1$, which demonstrated with alternation of X cm thick light greiwacke-rich interbeds with more pelitic (rich in mica) beds. The multi-stage of granitization process is distinct. Larger beds were formed in ductile stage of the thermal metamorphism of the rock, while thin veinlet (above hammer) is tied to semiductile flexure, originating in higher part of the crust (sample LHS-42).



Fig. 17. Crossing off the older granite position during ductile deformation, resulting in origin of oblique pygmatitic vein of granite (sample LHS-57)


irregularly dispersed in thin section. Rare there is present Chl in association with Grt. Pl has intermediate oligoclase composition (An_{23}), biotite and chlorite are probable coexisting minerals with roughly similar share of Fe/Mg. Grt is prograde zonal in the center with a predominance of the spessartite component over the almandine component, towards the edge with a predominance of the almandine molecule, a decrease in the grossular component and an increase in the pyrope molecule.

Chemical dating of monazite


Chemical dating of monazite from the sample LHS-350 (Plt. 1-1) provided identical result as Mnz dating from samples of further paragneisses. Distribution and age diagram of chemically dated monazites (Plt. 1-1) and isochrone with the age of 366 ± 6.8 Ma are typical for metamorphites of the Suchý hill. The data can be understood as a bimodal distribution corresponding to the




A




B



C



D



E

Fig. 18. A – Leucocratic melt at the boundary of granitoid part formed by hypidiomorphic plagioclase and quartz, containing relic biotite. Microphoto: melanosome composed of biotite and muscovite (after sillimanite) and chlorite. B–C – Detail melanosome formed of biotite and sillimanite N//, NX. D–E – Granite part formed of hypidiomorphic Plg and quartz. N//, NX (sample LHS-114A).




Fig. 19. BSEI of sample LHS-114A. A – Granite part – hypidiomorphic Pl without compositional zonation (An23-24). B – In granite part occurs biotite with rim of Ms, Ep and Kfs in plagioclase, phyllosilicates in granite part are similar regarding their composition as in the gneiss part. C – Melanosome is formed in a part of thin section by association Bt–Chl–Ms. Lamellae of ilmenite are common, being tied exclusively to muscovite and chlorite, which proves the later decomposition of biotite

peak of metamorphism around **380–370 Ma** and thermal reworking by granitization event **360–350 Ma**.

We have shown more significant beds of black graphitic paragneisses, black metaquartzites, which are in clear association with amphibolite bodies, in the area of Závada and Železná dolina valleys and in the area of Závadská poľana towards the Čierna Lehota village. Impregnations of sulfidic minerals are typical for this group of rocks, which create light, graphite-rich cavernous rock during weathering. In the environment of graphitic paragneisses and black metaquartzites, there are frequent traces of field exploration works. In the past in the area of the Závada valley, a forecast area was set aside for graphite raw material.

Lithotectonic unit consisting of a complex of proximal sedimentary facies of the continental slope of the Paleozoic sedimentary basin

This complex is built mainly by metamorphosed, flyschoid sediments of a sandy-loamy composition with a very small proportion of clayey-sandy facies and a very

small representation of basic metavolcanites, or even their absence. Depending on the depth of the erosional cut, these lithologies are strongly metamorphosed.

Migmatitized paragneisses locally with stromatitic and pygmatitic bodies of leucogranites (116)

The migmatitized paragneisses represent a part of the lower tectonic unit (it contains mainly metasediments of greywacke nature, which we affiliate with flyschoid sediments of the continental slope with the absence of amphibolite lithologies. They occur mainly in contact with hybrid granodiorites, with which they finger-like alternate and occur in a tight overburden of migmatites. They typically occur in the Suchý massif in the area of the Bystrica valley closer to Rudnianska Lehota municipality, or in the area of the Krstenica stream (Liešťanská dolina valley) W of Liešťany village. These underwent initial partial melting and are a suitable lithology for tracing deformation structures in relation to granitoid formation.

Within this rock assemblage, the gneiss component significantly predominates over the leucogranite compo-

ment, which is characterized by ductile granite structures manifested by bulging of granitoid material, which undoubtedly originated at different crustal levels (e.g. Figs. 16 and 17). Also from this example, it is possible to deduce the assumption that the main granitization process was more related to the contribution of heat from the intruding large granitoid masses, which is manifested by the different intensity of the migmatization process, than to deep crustal partial anatexis. At the same time, it is possible to state that the crustal thickening and melting in the deeper parts of the crust was followed by gradual decompression, the subsequent ascent of granitoid magmas into the metamorphic assemblage during the syntectonic regime and a change in the rheology of the metasediments from a ductile to a semi-ductile regime, and finally to a regime with signs

As is evident from the presence of leucogranite positions with melanosome rims, deeper crustal conditions of partial melting can be assumed through leucogranite pygmatitic structures, without significant interaction with the gneiss substrate, to semiductile structures with transversal tabular veins. The mentioned structures are a consequence of the decompression of the metamorphic complex. Larger bodies of hybrid (schlier) granodiorites that are associated with this complex could have formed as a product of segregation of granitoid material into suitable structures, but more likely represent independent pulses of granitoid magma into the metamorphic substrate.

Monazite dating from leucocratic melts, associating with migmatitized paragneisses and hybridic granitoids

Chemical dating of monazite indicates in high-metamorphic migmatitized paragneisses to presence of Mnz population, corresponding by age with pre-granite regional metamorphism, while all metamorphosed lithological members of the Suchý crystalline basement contain the Mnz population corresponding by age, where the chemical age approaches approximately **370–380 Ma**, which corresponds with Upper Devonian. Origin of first melts relates with later thermal overprint with the age around **350–360 Ma**, which corresponds with the boundary of Upper Devonian / Mississippian. First leucocratic melts contain mixed population of monazites from older metamorphic phase as well as

younger granitization stage. Following bigger volumes of hybridic tonalites and granodiorites contain only population of granitic stage.

Higher metamorphosed biotite to muscovite-biotite paragneisses with garnet, medium-grained, often with oriented secondary biotite (locally with sillimanite and staurolite) –distinctly injected with leucocratic granite (115)

In geological map we separate complex of biotitic and two-mica paragneisses, very intensively injected by leucogranites, aplites and pegmatites. At the same time – in profiles the paragneisses are alternating with paragneisses and granitoid injections in distances of several meters and dekameters.

Part of these rocks do not contain such leucocratic injection, but the presence of larger oriented and thermally induced biotite blasts, usually associates with nearby bodies of leucogranites in the footwall. The distinguishing of a zone of paragneisses with blastesis of secondary, thermally induced biotite in conditions of dynamic emplacement of leucogranite magmas indicated the vicinity of larger bodies of leucogranites and pegmatites–aplites in nearby footwall. Locally paragneisses form larger xenoliths in leucogranites. The presence of pygmatitic structures of leucogranites is absent, or is very rare, which indicates the higher crustal level of these lithologies during intrusion of leucogranite melt.

Thermal effect of granitoid intrusions produces larger flakes of biotite of 2nd generation, either in foliation planes of metamorphites, or growth of biotite II is characteristic for the whole rock. Blasts of biotite II have linear orientation and their arrangement corresponds with orientation of larger biotite flakes in leucogranites. Porphyroblasts of biotite II (Bt II) with flakes large up to 2–4 mm are characteristic for this type of gneisses (Fig. 21). They usually contain secondary garnet of contact thermic origin, reaching dimensions up to ca 0.1–0.2 mm (Fig. 25)







Fig. 20. Position of higher thermally affected paragneisses with thermally induced biotite II (115).


Fig. 21. Qtz–Pl–Bt–Ms paragneiss with older sillimanite–fibrolite (prevalence of Qtz over Pl). Sillimanite phenocrysts overreach 2 mm. At margining the fibrolite is replaced by Ms. The graphite pigment is abundant. Large blast of rotated sillimanite from regionally metamorphic phase is in later stage in connection with intrusive leucogranitic thermic stage cataclased and fracture is filled with red-brown Bt, which has different position concerning the older Bt of subparallel metamorphic setting of two-mica paragneiss. A – N//, B – NX (sample LHS-197). Grey fine-grained paragneiss, overheated by thermic metamorphism in connection with deformation related to ingress of leucogranites. Synintrusive deformation is manifested by the formation of microfolds and formation of larger flakes of phyllosilicates. Metamorphic schistosity is subvertical of E–W direction.



A



B



C

Fig. 22. Sample of biotite paragneiss with thermal overprint (LHS-255) and origin of redbrown biotites and tiny garnet. A – N//, B – NX, C – BSEI of disseminated tiny Grt in feldspars and Bt. Tiny Grt (below 0.1 mm) associate with Bt. Bt has intermediary ratio $Mg/(Fet + Mg) = 0.45-0.44$. Grt has strong prevalence of almandine component. Composition of Pl is in this association in the range XAn 29.5–32 %. Regarding very small dimensions of Grt we can state their relatively homogeneous composition in the profile. Typical average composition of non-zonal Grt is Alm – 71 %, Spess – 16, Pyr – 8, Gross – 4.6 %. Sample by taken from the contact of paragneisses with aplite veins and granite.




Plate 1. Mnz dating from granite part (sample LHS-114AG – Plt 1-2) is formed by Mnz, crystallizing from the melt and relict, being a part of relict minerals, mainly micas and monazite enclosed in them. The file can be divided to two age groups with the ages **350 and 390 Ma**.




Plate 2. Mnz dating from gneissic part (Plt 1-3) contains exclusively older generation of Mnz (368 ± 5.7 Ma). This age is characteristic for all distinguished types of metamorphites and most likely represents peak of regional metamorphism around 380–370 Ma.




Plate 3: Mnz dating from hybridic homogenous leucotonalite-leucotondhjemite (sample LHS-114B) (Plt 1-4), as ductile melt from the environment of migmatitized paragneisses, indicating the prevalence of Mnz age generation 350 ± 4.8 Ma.




Plate 4. Mnz dating from the sample of omnidirectional granodiorite LHS-94B providing the age 355 ± 5.5 Ma

It is proved by concordance of the main component of the stress field of synintrusive deformation regime with emplacement of leucogranite veins. These simultaneously with emplacement and decompression of the whole crystalline block rapidly solidified and behaved as dense crystal slurry.

They have corresponding mineral composition as following type of paragneisses, but slightly larger grain. In prevailing lepidogranoblastic structure the bands of biotite-plagioclase material alternate with cleaner quartz-rich lamina. Plagioclase grains usually connect by polygonal way and obtain hornfels structure. In comparison with the standard biotite gneisses, gneisses of this type were to some extent injected by leucocratic material. In the process of “thermic” metamorphic crystallization they acquired larger grain size, and probable there rarely crystallized also transversal muscovite.

Chemical dating of monazite in metapsammitic paragneisses

Monazite in biotitic paragneisses is relatively abundant mineral, often containing according the results of dating

at least two separate populations. In paragneiss sample LHS-255 two temporal groups of monazite occur. Older group corresponds to boundary of Silurian / Devonian (410 Ma) and probable can be correlated with the period of filling sedimentary basin with clastic material? Age 380–370 Ma is significantly more distinct and probable relates with collision and metamorphic overprint in Upper Devonian. Following thermal metamorphism of Lower Carboniferous age was only indistinctly reflected by the origin of new generation of monazite (Plt 2-1).

Lithotectonic unit consisting of orthogneisses complex (granite protolith Cdd2, tectonometamorphic overprinting Vmd2)

Biotite, quartz-plagioclase (\pm K-feldspar \pm muscovite), foliated and lineated orthogneisses (111)

Quartz-, plagioclase-biotite orthogneisses with variable content of interstitial K-feldspar and muscovite crops out mainly in separate, distinctly limited zone with sharp contact with neighbouring biotite paragneisses. Main body




Fig. 23. Complex of orthogneisses (111) forms supreme structural level of pre-granite structural arrangement. It represents shallow deposited fan. Manifestations of a limited degree of partial melting are developed only in its root part in the Nitrica river cuts, probably with the contribution of shear heat near the contact with the paragneiss complex.

of E–W direction is situated south of Valaská Belá village approximately W of the elevation point Capárka (924 m) through Okružly vrch (914 m) towards the confluence of the streams Jasenina and Nitrica (area at the crossing at the Motorest Klin). In this area the body of orthogneisses is areally reduced, which is caused by deeper erosion cut with higher degree of granitization, connected probable

with the contribution of shear heat. This fact indicates the upper structural position of the body within Variscan tectonic setting (profile 1–2 in published geological map) in the footwall with the paragneiss complex. In this zone we interpret this position as fan structure, wedging into the deeper parts. Further this body follows in E–W to ENE direction towards the area of fault east of Temešská




Fig. 24. A – Banded orthogneiss with transversal vein of leucocratic granite (sample LHS-207); B – ductile deformed relics of plagioclases in orthogneiss (sample LHS-273).




Fig. 25. Orthogneiss with plane-parallel setting, formed of chestnut-brown Bt (sample LHS-272.) In Qtz-feldspar matrix moderately prevails Qtz over Pl and K-feldspar. Ap, Zrn are frequent as accessory minerals, Mnz is present, too. The feldspars porphyroblasts in the rock occur in the form of quartz, plagioclase and Kfs aggregates.
Microphoto: A, B – N//, NX, C – BSEI of the orthogneiss structure.

skala hill. Here it is intruded by younger leucogranites and biotite I-type granodiorites (106).

In the past, this lithology was considered as migmatites. However, the processes of migmatization in the rock are completely absent and are limited only to narrow zones, even in the deeper parts of the fan-shaped structure.

These are monotonous banded rocks, where quartz-feldspar bands alternate with biotite bands a few mm thick. The content of biotite and plagioclase slightly varies. The presence of spindle-shaped plagioclase is typical. The metamorphic foliations are steep, mainly in the E–W direction, with lineations mainly of biotite glomeroblasts of a subhorizontal direction, which contrast with the light quartz-feldspar mass. The position of orthogneisses during granitization processes was more external (higher), which is indicated by only very rare ductile deformations of orthogneisses associated with the presence of leucocratic melts. The aplite and pegmatite veins present in them are usually oriented at a high angle to the foliation of the orthogneisses.

The dominating foliation features indicate the presence of ductile shear folds with steep fold planes, shallow fold axes, and shallow lineations accentuated by elongation of biotite glomeroblasts. The pre-deformation origin of orthogneisses is not clear. Unlike the lighter, fine-grained paragneisses, the rock lacks graphitic pigment, which shifts our opinion to the assumption that it is an acidic, originally magmatic rock of granitic to granodiorite composition.

Monazite chemical dating

Chemical dating of monazite from orthogneiss (sample LHS-711) indicates an average age of 360 Ma, which, however, does not correspond to a real geological event. Dividing the data into two distinct peaks – into two separate events, where we interpret the older event (400–380 Ma) as the age of metamorphic event. The second significant maximum is the age of 355–345 Ma, which corresponds to a younger thermal event associated with the formation of granitoids in the Strážovské vrchy area (Plt 2-2).

Migmatites with prevalence stromatitic, loc. ptygmatitic textures (117a)

Migmatites form the lowest structural level of the crystalline massif from the point of view of the Variscan (pre-granite) setting. They emerge in the SE edge of the Suchý massif (so-called Liešť migmatites). Pre-granitization, or syngranitizing fold planes and elements of metamorphic foliation have a steep to subvertical course in the NNE–SSW to NE–SW direction. The axes of the folds are predominantly subhorizontal.

In *stromatitic migmatites* there varies the content of the light quartz-feldspar component representing the neosome relative to the relict part, locally enriched with minerals rich in Fe and Mg (melanosome). The content of light and

dark components varies from types with a predominance of the light component to the dark component in a ratio of approximately 70 : 30, to types with a slight predominance of the dark component over quartz-feldspar component. It points more to the variability of the composition of the substrate than to the intensity of the anatexis process. The process of anatexis led to a maximum of the first few tens of percent of the share of the original volume of the rock. Smaller volumes of migmatized pararules are also present, where, in addition to granitoid injections, more or less numerous ptygmatitic granitoid melts appear.

In the Suchý massif, a strip of migmatites is adjacent from the western side to lighter schist-like (hybrid) granodiorites (110), which represent an overlying complex to the migmatites. Although the contact is relatively sharp, larger migmatite septa are found parallel to the foliation elements in the migmatites and parallel to the inhomogeneities – schliers in granodiorite. Among the migmatites and hybrid granodiorites, amphibolite bodies that have undergone granitization occur. From a lithological point of view, this type of migmatites may correspond to the original composition of the orthogneiss material.

The composition of large flakes of biotite, characterized by a stable composition with $Mg\# = 0.38–0.40$, is close to the composition of muscovite with $Mg\# = 0.43$, which points to equilibrium conditions of association. Ms is often very intensively replaced by sillimanite by the reaction $Ms + Qtz$ to form a melt, resp. $Bt + Qtz$ to form a K-feldspar-rich melt (outside of the presented thin sections). Putiš (1976) reports the volume composition of mineral components from the more isotropic type of migmatite SW from Temeš village – quartz (19.2 vol. %), plagioclase-oligoclase (27.9 %), orthoclase-microcline (19.4 %), biotite (18.1 %), muscovite (2.5 %) and sillimanite (12.4 %).



Fig. 26. Ductile deformed migmatites with equal representation of light and dark component. Right beneath the hammer the bounds of quartz-feldspar pegmatite-type melt are visible (sample LHS-131 – the Krstenica valley). Structure of the rock is grano-lepidoblastic to lepidoblastic with dominating plagioclase, biotite, K-feldspar, muscovite, sillimanite.




Fig. 27. Mineral association of melanosome from migmatite is formed in prevalence of biotite associated with Ms, Qtz and Pl replacement by Bt and Ms by sillimanite left part of the picture (sample LHS-137), BSEI.




Fig. 28. Association Ms + Bt melanosome (apatite inclusion). Black subparallel tables of graphite enclosed mainly in Ms. Subparallel arrangement of graphite strips shows that this part represents former metamorphic assemblage. Chaotic arrangement of graphite right down points on flow arrangement due to melting of former consumed Bt (light relic right down). Ilmenite with content up to 5.61 wt.% of MnO crops as akcesoric mineral together with plagioclase with An₁₈. Ilmenite and graphite are oriented according to metamorphic foliation surfaces (sample LHS-137). BSEI.

Late Devonian – Mississippian granitoids (Vmd2)

In the Suchý massif there is possible to distinguish several granitoid portions, which are close in time, but intruded at specific crustal conditions in decompression regime. In lower part of metamorphic packet there intruded the schlier granodiorites with less mobile magma, which is proved by inhomogenities parallel with those

developed in migmatites. In schliered shallower crustal conditions the transtension shearing applied with intrusions of S-type prevailingly two-mica granites (connected with distinct presence of transtension pegmatite intrusions), up to intrusions of I-type granodiorites, or even small diorite bodies.

Massive to weakly oriented hybridic granodiorites with common schliers and xenoliths of paragneisses and migmatites (110)

Schliered granodiorites emerge to the west of the migmatite zone, while their mutual relationship is relatively sharp, but blocks of underlying migmatites can be found quite commonly, especially in the contact zone, about a few more 100 m from the mapped border with migmatites. The contact is highlighted in places by smaller or larger bodies of amphibolites. The higher part of the complex is dominated by bodies of migmatized paragneisses and biotitic to two-mica paragneisses. These granitoids crop out in the zone of NNW direction from the Bystrica valley W of Rudnianska Lehota, where they alternate with migmaritized paragneisses (116), through the localities Háj – Predné lazy – Prostredná dolina – Obory nad dolinou Krstenica




Fig. 29. Position of schliered granodiorites (110) in the lower part of crystalline pile.

(Liešťanská dolina valley) – Mihálová (where they are in contact with omnidirectional younger biotite granodiorite) – towards Nitrica and in the E this zone ends on fault zone of N–S direction.

The preferred orientation of foliation structural elements of NNE–SSW direction in migmatite and paragneisses blocks is identical and corresponds with preferred orientation of dark biotite schliers in granodiorites.

Granodiorites represent medium-grained granitoids, where the quartz-feldspar component is more or less omnidirectional and the inhomogeneities are formed by biotite schliers, gneiss and migmatites xenoliths, which setting indicates preferred orientation. However, there are also present varieties with a low content of inhomogeneities and omnidirectional biotite.

Xenoliths of paragneisses and migmatites range in size from a few cm to hundreds of meters in places. The composition of granitoid rocks varies from tonalite to granodiorite.

Schliered biotite granodiorite–tonalite (sample LHS-177-2) contains moderately zonal plagioclases of composition An_{24-28} and biotite with $Mg \# = 0.41-0.44$. Rare Ms appears in the structure as later, which is indicated by different $Mg \# = 0.52$. Zircon, monazite, apatite and ilmenite are rare present as accessory minerals.

Mnz dating based on 27 values gave a fairly uniform distribution, where the age $348 \pm 4,3 \text{ Ma}$ (sample LHS-177-2, Plt 2-3) can be considered as an age close to the thermal event that led to the emergence of melting processes and subsequent crystallization.

Schliered granodiorites represented the less mobile crystal slurry, which is associated with the underlying migmatites and was generated by ongoing thermal reworking of the metasediments. The rock is essentially a diatexite, i.e. a rock that has passed through the initial magmatic stage, which is evidenced by the zonality



Fig. 30. A – Schliered granodiorite with xenoliths of migmatized paragneisses and oriented biotite.



Fig. 30. B – A schliered granodiorite with a substantial representation of an inhomogeneously distributed components, subparallel, unevenly distributed laths and accumulations of biotite, which represent a redistributed melanosome. In some places, the biotite melanosome forms separate areas of biotite, separated by fuzzy pale stripes: The rock represents a higher degree of melting and mixing of granodiorite leucosome with melanosome in the form of a dense crystal slurry (sample LHS-185).





Fig. 31. Structure of the schliered granodiorite with oriented Bt laths, xenomorphic to hypidiomorphic weakly zona Plg. Ms is rare and appears in the structure as late in origin (sample LHS-177-2).

of plagioclase and biotite orientation, the presence of numerous schists, migmatite xenoliths and migmatized paragneisses. The separation of the K-feldspar component is only very limited and manifests itself in isolated positions parallel to the inhomogeneities in K-feldspar rich grey parts. Despite the rheological properties, the mobility of the crystal slurry can be observed in places, which is manifested by sharp contacts with the migmatites, which appear as septa in this granitoid.


Biotite or muscovite-biotite granites, locally leucocratic (109)

They form a strip emerging west of the rock type No. 110 and are spatially and genetically connected to it. They represent a band of granitoids dominated by more leucocratic types of granodiorites. They have a lower content of dark components.


The rocks have a slightly oriented biotite, which is generally inhomogeneous and size-distributed. But




Fig. 32. The character of medium-coarse-grained, more leucocratic types of granodiorite rocks with irregular spatially and size-distributed biotite.




A



B



C



D

Fig. 33. Microphoto of the rock type in Fig. 32 – association Bt–Grt–Ms from the schlier, B – muscovitization of sillimanite, C – association quartz–plagioclase–Bt–Sill (gray tufted objects), D – replacement of Bt in association with sillimanite into the quartz-feldspar matrix by fan-shaped Ms. A, C = N//, B, D = NX (sample LHS-94 A).




Fig 34. BSEI of biotite schlier (A – general view, B – detail). Kfs (light grey phase in Fig. B upper left) due to its morphology, it is a product of the melting reaction of biotite to form sillimanite + K-feldspar and garnet. The lightest phase represents a relict of decaying biotite with overgrown sillimanite, and the slightly grey phase is decaying muscovite with inclusions of euhedral sillimanite. The second stage, in subsolid conditions, is the replacement of the original association by muscovite, in which relics of sillimanite are still preserved (upper left under Kfs).

there are also varieties with omnidirectional Bt. The Bt dimensions sometimes exceed 1 cm. The different size and spatial characteristics of Bt indicate that only a small amount of the biotite crystallized from the magmas. They contain significantly fewer xenoliths of paragneisses as well as biotite schists compared to lithotype No. 110.

The structure is dominated by plagioclase (up to 7–8 mm in size), which is hypidiomorphic to xenomorphic, usually oriented in the direction of magmatic flow (or flow of crystal slurry), which is simultaneously parallel to the course of isolated schliers. Interstitial Kfs has a relatively small presence. Rarely, Grt is macroscopically observable together with Bt. Sillimanite was observable only in thin sections. Grt is moderately zonal with a retrograde type of zonality from the center (Alm 74 – Pyr 12.3 – Spess 10.7 – Gross 3.0) towards the margin (Alm 73.1 – Pyr 8.5 – Spess. 15.2 – Gross 3.2). Biotite has a prevalence of Fe – Mg# = 0.38–0.40.

Leucocratic two-mica granites (107, 108)

This type of granite is dominantly developed in the central zone of the Suchý massif. These subvertical intrusions are associated with a process of shear deformation with a NE–SW direction. The importance of this deformation is manifested in the surrounding metasediments by opening of spaces for linear types of intrusions, in connection with their thermal effect. Shear deformation occurred continuously from the suprasolidus to the subsolidus stage, which is consistent with the scenario of decompression emplacement of granitoid intrusions.

Leucocratic syn-intrusive oriented two-mica granites (108)

Leucocratic two-mica granites are the most widespread petrographic type in the Suchý massif, building mainly its central part. They are spatially connected mainly with paragneisses (115), into which they intensively penetrate. However, they also form separate stocks and linearly oriented strips, preferably trending NE–SW. From the point of view of the Variscan setting, they represent the upper part of a complex intrusive body, while the content of muscovite in the granitoids increases in the direction towards the overburden, i.e. to the NW. From the point of view of the genesis of pegmatites, it can be stated that only in these leucocratic granites can one observe gradual transitions from zonal bodies formed on the edge by aplite and in the central part by pegmatites with an albite fine-grained zone and the central part formed by blocky grey K-feldspar, quartz with occasional muscovite and biotite. Ms appears in the texture as primary, but also as later one, which is evidenced by the growth of Ms at the expense of sillimanite.

It follows from the above stated that the majority of pegmatites can be associated with leucogranites of this type, where in the contraction zones formed at a steep angle to the linear flow of leucogranite magma, cracks were formed in the high-temperature regime, being filled with pegmatite melt together with enrichment with a fluid component. The course of the pegmatite veins is dominantly N–S with variations the NNW or NNE direction.




Fig. 35. Different types of planar oriented leucogranites: A – Planar oriented two-mica granite in the subsolid stage, with a pegmatite vein without deformation (sample LHS-626). B – Planar oriented two-mica granite up to developed the subsolid quartz bands (sample LHS-623).





Fig. 36. A stock of muscovite-biotite granite that intrudes into biotite paragneisses (about 100 m from the contact with them). The granites are fine- / coarse-grained leucocratic, sometimes containing Grt. Pegmatitoid granites with Grt are also common. Omnidirectional leucogranite with a predominance of Ms over Bt. It contains a microcline that poikilitically encloses oval Qtz and Pl I (up to 0.2 mm in size; Fig. B). Fibrolitic sillimanites are present, being replaced by large flakes of Ms. But there is also present longer prismatic Na-richer Pl, which has a Ms shell in the vicinity without fibrolite. Pl encloses the oval Qtz I, which, due to the high content of Si in the melt, crystallized among the first minerals. Pl is composed of albite to Na-oligoclase (X_{An} 3–16) and is only indistinctly zonal, Kfs contains 7.6–10.9 vol. % of albite molecule. Bt is slightly directional and is probably relict. Grt was not revealed in the thin section, tiny pink Grt is macroscopically rarely present. The average grain size of the rock is 1–2 mm. A – texture of two-mica granite, where K-feldspar encloses Pl and Qtz of the 1st generation; B – An omnidirectional microcline closing Qtz I and Pl I; C – Fibrolitic sillimanite replaced by muscovite. D – Long-prismatic Pl, rimmed with Ms flakes, partially closing Qtz I. *NX* (sample LHS-73).

The flow structure resulting from the magmatic flow direction is finally reflected from the early stages of crystallization up to the subsolid stage. It corresponds to the decompression stage of magma emplacement as well as the entire crystalline complex. The emplacement of these leucogranites took place partly even in the subsolid regime, while the foliated types of leucogranitoids were formed, which can be found in the footwall of injected gneiss complex (115) in the source area of the Krstenica stream (Liešťanská dolina valley). At the same time, the dynamic placement of leucogranites leads in the adjacent gneiss lithologies to the origin of preferred orientation, rotation of the former blasts of regional-metamorphic minerals (e.g. sillimanite, staurolite), the formation of


larger linedated blasts of biotite of the second generation (Bt2).

The leucogranites are mostly medium-grained, with a small representation of Bt. Their texture varies from omnidirectional to that with preferred orientation of newly formed minerals (linedated Bt) but also planar foliated. The structure is hypidiomorphically granular. Rocks have a multi-stage crystallization history of rock-forming components.

Based on microprobe analyses, the plagioclase composition ranges from An₂ to An₁₇, K-feldspar – with max. content of albite component – 10 vol. %. Muscovite has content TiO₂ 0.15–0.75, which points to the fact that it probably did not crystallize within the magmatic stage, but




A




B

Fig. 37. Muscovite-biotitic (Ms : Bt ratio about 1 : 3) equigranular omnidirectional granite. Bt and Ms, as well as Pl ± Qtz are moderately oriented, while the small Qtz and Kfs are without preferred orientation – omnidirectional. The myrmekites are developed in places among Kfs and acidic Pl. This indicates synintrusive deformation due to flow direction. The average grain size of the Qtz-feldspar material is 1–2 mm, the average size of flakes of Ms up to 0.5 mm and Bt up to 0.1x0.4 mm (sample LHS-200).



A



B

Fig. 38. Leucogranite. A – Sillimanite with Bt relicts is replaced by muscovite, N//. B – Manifestation of subsolid deformation in the solid state (zone in the middle of the image - red arrow), which is nearly parallel to the earlier crystallizing Pl (sample LHS-206).

is younger and partly formed at the expense of sillimanite and biotite. The rarely present garnet is rich in almandine component (Alm – 78, Pyr – 8.7–10.3, Spess – 9.2–11.5, Gross – 2.3–2.4). The Hg content of biotite is low – $M\# = 0.31–0.39$, muscovite – 0.40–0.47. Content of TiO_2 in biotite varies from 2.2 to 3 wt. %.

Leucocratic omnidirectional to weakly oriented two-mica granites (107)

They occur together with the previous type of more syndeformation oriented granitoids. Similar to the previous type of leucogranites, the deformation is caused by the magmatic flow, concordantly with the shear deformation in the surrounding metamorphites.

Dating of monazite from two-mica granites indicates a relatively narrow intrusive age range of **352–351 Ma** (sample LHS-73, Plt 2-4; sample LHS-482B, Plt 3-1 and sample LHS-482A1, Plt 3-2), which can be considered as the age of maximum granite forming process (intrusive phase in the shear regime). At the same time, the mentioned granites are characterized by the rapid dynamics of crystallization of mineral components. If the initial phases (crystallization of mainly plagioclase and coexisting micas) are oriented in the magmatic flow, the final phases are already more or less omnidirectional. However, already rheologically solidified rocks are still further deformed with the origin of planar structures in places (Fig. 35).

The whole process is thus related with the ongoing deformation of the rock complex and at the same time the decompression of the entire set of metamorphic and igneous rocks (rapid ascent to higher levels of the crust). Residual melts of aplites and pegmatites, associated with this complex, are bound to ongoing deformation in the semiductile to brittle stage of magma solidification and are placed in extensional structures that arise at a large angle to the main component of tectonic stress.

Pegmatitic and aplitic veins (103)

Pegmatites and aplites are genetically related to the development of the shear system and intrusions of two-mica leucogranites. As a rule, they fill transversally oriented brittle structures, oriented at a large angle to the main stress component. In the Suchý massif, their course is usually oriented N–S (varying within NNE–NNW).

Pegmatites and aplites are genetically related, with aplites forming the external part of a complex pegmatite-aplite system. The marginal, aplite part is made up of fine-grained “sugary” aplite, whose mineral components, especially quartz and feldspar, are oriented omnidirectional, unlike it is in leucogranites. The macroscopic pink garnet that is often present in the aplite structure is striking, and can range from mm to several cm in size. Although

the aplites crystallized together with the pegmatites as a final stage, some orientation caused by magmatic flow in a dynamic environment is evident from the occasional linear arrangement of garnet phenocrysts (Fig. 39D). Pegmatites are coarse-grained and formed from the edge by a less coarse-grained zone with a predominance of albite feldspar, while inside they are formed by blocky quartz and grey K-feldspar, which sometimes reaches several dm to m in size. As a rule, pegmatites contain little mica, but there are varieties with large-scale Ms and, in places, Bt, which is preferably bound to the marginal part of the pegmatite bodies. The relation to the metamorphic rocks and older granitoids is sharp, and the aplite and pegmatite bodies occur in the form of sharply bounded tabular bodies. The only exception is the relationship to two-mica leucogranites, where in some places indistinct transitions from leucogranite and aplite to pegmatite are observable. This fact points to the genetic relationship of these lithologies.

The abundance of aplite and pegmatite veins is highest in the central part of the territory SE, ENE of Suchý vrch hill, while it is mainly linked to leucogranites and the overlying paragneiss complex, which is in contact with leucogranites. Injections of leucogranites and pegmatites – aplites often alternate in the paragneiss complex (115). In the direction to the SE – into the underlying formations, the number of pegmatite and aplite veins decreases, and the same occurs in the direction to the NW and to marginal (upper) parts of the crystalline basement. Pegmatite and aplite bodies are not always complexly developed in the zonality described above. In places, only separate swarms of aplites are present, or only swarms of pegmatites.

Regarding the cartographic visualization in the published geological map (Hraško & Kováčik, eds., 2021; web link available), from the genetic point of view we do not differentiate between pegmatites and aplites. The course of the bodies in the geological map does not always correspond to the actual course of the veins, but represents the concentration of vein swarms in the given area.

According to the chemical dating of the monazite, as well as according to the geological position, these bodies represent younger magmatic stage of the shear emplacement of the granitoid magma – **345–343 Ma** (sample LHS-557, Plt 3-3; sample LHS-607, Plt 3-4).


Aplite garnets are almost non-zonal with high almandine content, in association with Fe-rich Bt with $M/MF = 0.35–0.36$.

Chemical dating of monazite from two selected aplite samples indicates consistently low ages in the given range of Lower Carboniferous ages. Age **345 ± 2.9 Ma** up to **343 ± 2.9** can be reliably considered as the conclusion of magmatic processes associated with the shear emplacement of leucogranites and aplite-pegmatites.



- A – Up to 3 cm thick transversal pegmatite veinlet with grey Kfs, forming tabular body with oblique orientation of ca 75° to course of lineation of biotite glomeroblasts biotitu in biotitic orthogneiss (sample LHS-415).
- B – Wedging of the Ms pegmatite vein in Al-rich (metapelitic) paragneiss (Stau-Sill), at the time of intrusion the paragneiss was partially plastic (sample LHS-555).

- C – Several cm large garnet in coarse-grained quartz-albite part of pegmatite (sample LHS.599).
- D – Omnidirectional “sugar” aplite with small pink Grt on the edge of the Ms pegmatite (sample LHS-572).



- E – Partially weathered pegmatite, where prominent blocks of grey K-feldspars are chaotically arranged in quartz-albite fine-grained mass. The texture is a result of subsolid (pneumatolytic?) fracturing (sample LHS-611).

Fig. 39. Typical present surface position of pegmatite and aplite bodies:

From the S-type highly orogenic granitoid stage to the I-type late-kinematic granitoids

A special type of I-type granitoids mainly builds up the NW edge of the crystalline area, to a limited extent in the Suchý massif, but especially the eastern area of the Magura part chemical basement. From the point of view of using dating of monazite, it was not possible to decide on the temporal relationship of the leucogranite and aplite-pegmatite intrusive stage (350–345 Ma) associated with the shear-deformation regime and the intrusion stage of I-type magmas. However, from the point of view of spatial relations, it is clear that I-type intrusions have a late kinematic character in relation to the shear deformation of the complexes.

Biotitic granodiorites – tonalites, omnidirectional, only locally weakly oriented (106)

Biotite granodiorites form several separate bodies of predominantly isometric shape, which points to their later age compared to the shear-deformation events of the Lower Mississippian. They mainly penetrate these structures. Smaller bodies emerge on the E-oriented slope of the Suchý massif, W from Nitrica, and the larger body emerges in the area of Čavoj locality.

Granodiorites are conspicuous by their omnidirectional structure, fresh appearance and the lack, or the complete absence of pegmatite veins. This fact also points to their younger and deeper magmatic portion in the Variscan geological structure of the crystalline basement. In some places, several more mafic “globules” of omnidirectional granodiorites to tonalites can be observed, which interpenetrate each other in the form of loaf-like bodies of




Fig. 40. Late kinematic intrusion of I-type granodiorite (106).

different grain sizes (Fig. 41). This indicates a good viscosity and a higher temperature of the magma.



Fig. 41. Biotitic granodiorites – tonalites – mixing of two types – block of more mafic part rich in plagioclase and the biotite rich part (above hammer) in more quartzy coarser-grained rock.

From a petrographic point of view, rocks represent biotitic granodiorites to tonalites. The Pl (up to about 50 %) rich granodiorite, which is also rich in Bt (up to 20 %), while in some places it may represent portions of mafic magma in a lighter type (Fig. 41). Pl usually 1 mm (rarely 2 mm) large are idiomorphic to hypidiomorphic, the intergranular spaces are filled mainly with Bt and Qtz.

Hypidiomorphic plagioclase is zonal, oriented in all directions, without inclusions, only in the final phase it crystallized together with quartz. Biotite with a



Fig. 42. Fine-grained omnidirectional, grey biotite-rich microgranodiorite-microtonalite, as one part of the biotite granodiorite-tonalite body (sample LHS-122 – Krstenica valley).

content of 15 to 20 vol. % is reddish-brown in colour and appears usually along the edge of plagioclase, from the point of view of succession it is later than Pl. As a rule, quartz and quartz are interstitial and crystallized last in the succession. Of the accessory minerals, apatite (Ap) and also monazite (Mnz) are commonly present, which were chemically dated here. Magnetite is also present, which indicates a higher activity of water in the magma, as it points to an oxidation regime in the magma, which was a consequence of the dissociation of water vapor in the magmatic reservoir.



Fig. 43. Granite sample LHS-94B. Hypidomorphic plagioclases are zonal, crystallizing first in crystallization succession. Only rarely in their final phase of crystallization the interstitial K-feldspar crystallized. NX.

Mineralogy: Plagioclase, K-feldspar, biotite and rare muscovite were measured in the granite in association with Bt. The composition of plagioclase measured by microprobe analyser is An26 to An36. The content of the albite component in K-feldspar reaches 10 vol. %. the BaO content is around 2.5 wt. %. The magnesium content of biotite (Mg#) varies in a narrow range of 0.51–0.52 and the TiO₂ content – 3.4–3.65 wt. %. The Mg content of Ms is around 0.57.

Chemical dating of monazite (Mnz) from a sample of omnidirectional granodiorite (sample LHS-94B, Plt 4-1) provided an age of **355 ± 5.5 Ma**, while the statistical distribution of the set points to one, homogeneous population of monazite.

Coarse-grained amphibole diorite (125a)

Such coarse-grained, to medium-grained omnidirectional, to weakly oriented amphibole-plagioclase rock was found in the environment of the previous type of omnidirectional granodiorites-tonalites of the I-type (sample only in scree).

The amphiboles correspond to the composition of magnesiohornblende, the plagioclase that fills the spaces between the amphiboles is of andesine composition. The association of ilmenite and titanomagnetite and then rutile and titanite is common, which indicates the oxidation regime in the magma and sulphide minerals (pyrite, chalcopyrite?) are present. The stated oxidation conditions correspond to the conditions of magmatic crystallization of the intermediate magma. Therefore, we consider this lithology to be a part of the previous type of granodiorites to tonalites with features close to I-type granitoids, with higher temperature and water-rich magma.




Fig. 44. BSEI of sample SZN-304 consisting of phenocrysts of magnesiohornblende and interstitial andesine. Former ilmenite is replaced by rutile and titanite aggregate due to postmagmatic oxidation.

Discussion and Conclusion

Character of pre-granite lithotectonic units present in crystalline basement of the Suchý massif

Pre-Alpine crystalline basement is built of variegated Paleozoic rock sequences regarding their position and depth in sedimentary basin, where they originated before they were amalgamated by Variscan tectonometamorphism process:

- Paleozoic deep water euxinic facies, formed originally mainly by fine-grained pelitic and quartzitic sedimentary facies rich in organic matter, with preserved fragments of accompanying oceanic basic volcanites. At field research these associations are relatively well detectable;
- Proximal sedimentary facies of continental slope of Paleozoic sedimentary basin, represented mainly by flyschoid sandy-greywacke sediments. They they contain almost no bodies of basic rocks and pelitic lithologies are very rare;


- The upper part of present Variscan setting is built of orthogneiss complex (less in W part of Magura massif and more extensive in the Suchý massif). In Variscan collision the orthogneiss complex as crustal granitoid segment was thrust on two above described rock sequences and later metamorphosed together with them.

Relations of metamorphic complexes and granitoid intrusions

Due to the Alpine tectonic rotation of the blocks of crystalline basement and cover Mesozoic sequences, the Suchý massif provides a more comprehensive profile through the basement, indicating an increase of regional

overprint from NW to SE. The highest degree of ductile deformation reworking is present in stromatitic migmatites, which typically crop out in the valley of the Krstenica stream (NW from Liešťany municipality), up to the area of Rudnianska Lehota. Biotite dehydration melting processes were also observed here. At the same time, intense granitization processes are present in this SE part, which are getting weaker towards the W–NW. For this reason, there were analysed the p-T metamorphic conditions from the W part of the crystalline basement, with minimal presence of granitization processes. On the other hand, to determine the pressure-temperature conditions due to the influence of intruding granitoids, samples were selected from the area of intense manifestations of granitizing activity.

Model of lithostratigraphic and lithotectonic Palaeozoic development of the crystalline basement of the Strážovské vrchy Mts.



Ages: Internat. Commission on Stratigraphy v 2022/10

Fig. 45. Lithostratigraphic and lithotectonic column with indicated position of protolith in sedimentary basins of various crustal provenience, their following amalgamation at collision, metamorphic overprint and syn-kinematic up to late-kinematic granite forming phase. Subsequent post-kinematic unroofing relates with orogenic collapse and ascent of crystalline basement to higher crustal levels (used ages are from actual International chronostratigraphic chart). OG – granitoid orthogneisses, gpe – sediments with predominance of pelitic component, gps – sediments with predominance of psammitic component, A – amphibolites-metabasalts, U – ultramafites, qg – quartz sediments with organic matter (black metaquartzites), C – metamorphosed carbonates, sM – stromatitic migmatites, GDh – hybridic granodiorites, IG – leucogranites, GD – biotitic omnidirectional granodiorites, P – pegmatites.

Regional metamorphic conditions correspond to temperatures of about 570–620 °C at pressures of about 6–8 kbar. Conversely, mineral associations of younger, contrasting thermal metamorphism correspond to approximately temperatures of 550–600 °C and pressure of 3–4 kbar (p-T calculations of metamorphic events were carried out by R. Demko, ŠGÚDŠ, Slovakia).

These different p-T conditions indicate a rapid uplift of the complex at the simultaneous intrusion of leucocratic granites in a shear regime. Due to the consistent directional relationship of deformation structures in leucocratic granitoids and surrounding metamorphites (concordant lineations of biotite schliers in leucogranites and thermally induced flakes of biotite II in the surrounding metamorphites), it is obvious that the leucocratic granites were emplaced simultaneously with the deformation of the metamorphites.

At the same time, it is obvious that the emplacement of leucocratic granites also took place in the subsolid stage, which is manifested in field conditions by the plate disintegration of leucogranites and deformation microstructures. In the subsolid stage, pegmatite bodies were also placed in the same deformation regime. Their composition (lack of water-bearing minerals) as well as textures indicate open structures and leakage of fluids into the higher parts of the metamorphic packet.

The temperature of regional metamorphism in the western part of the territory (the Suchý massif) did not generally reach the temperature of partial melting, but the granitization manifestations were mainly caused by the combination of the heat supply of granitoid masses in the peak stage of the syncollision event with simultaneous deformation. In the eastern part of the Strážovské vrchy Mts (Magura massif) crystalline basement, the processes of migmatitization were more distinct, because this part represented a deeper structural segment.

Tab. 1

Determination of petrogenetic parameters according White et al. (2014) and applying garnet-biotite-plagioclase-quartz thermobarometry (GBPG) after Wu, Chun-Ming et al. (2004)

Sample	Temperature (°C)	Pressure (kbar)	Comment
M2 granite induced thermal overprint			
LHS-204	593 ± 10	2.9 ± 0.2	thermal decomposition of muscovite to Kfs
LHS-356a	571 ± 13	3.4 ± 0.3	Bt, Ms, St, Grt (inclusion in St), Pl, Qtz
LHS-356b	578 ± 4	3.4 ± 0.1	
SZN-34a	580 ± 9	3.6 ± 0.3	Grt as inclusion in St, Sill and Kfs (product of high thermal decomposition)
LHS-329	561 ± 14	3.7 ± 0.7	cellular, poikilitic garnet, towards rim
LHS-255	585 ± 6	4.0 ± 0.2	enormous thermal blastesis o small Grt crystals
SZN-34c	591 ± 8	4.2 ± 0.2	
M 1 regional metamorphism preceeding granitic intrusions			
LHS-350	561	5.4	
LHS-329	581 ± 5	5.6 ± 0.5 ?	cellular, poikilitic garnet, central part
SZN-34b	582 ± 2	6.7 ± 0.1 ?	
SZN-66	608 ± 15	7.5 ± 0.8 ?	garnet I, in the centre of garnet II
LHS-350	569	8.5	




Fig. 46. Presentation of conditions of Variscan regional metamorphism (M1 reg.) in the upper part of the metamorphic pile (blue arrow). Melting reactions must have taken place mainly with the participation of muscovite melting. However, the speed of the process was not sufficient for more extensive melting and segregation of granitic magma. Granitic magma was generated from deeper parts under the influence of new thermal input (red arrow) and caused (M2 gr.) thermal metamorphism in higher crustal levels. The pink rectangle shows the near solid placement of the leucocratic magmas. (field of water-saturated granitic solid, mica melting fields and amounts of water provided during melting processes in metasediments according to Vielzeuf & Montel, 1994).

Deformation and magmatic structures in granitoids and surrounding metamorphites

The oldest Variscan structures are observable only in a microscale as oriented blasts of quartz within younger and larger blasts, especially staurolite. This oldest foliation is designated as sV1. The blastesis of staurolite originated in metapelitic assemblages at the peak stage of orogenic (regional) metamorphism, before the beginning of a younger deformation event, which caused the origin of the most prominent metamorphic foliation sV2 connected with shear deformation, thermal metamorphism under lower pressure conditions, being a reflection of granitoid intrusion in the shear-deformation event.

This younger deformation (sV2), which led to the rotation of older blasts together with their inclusions, is macroscopically and microscopically the best structurally

defined phenomenon. It led to the formation of steeply (up to subvertically) built structures of metamorphic schist, which were used in parallel syntectonic leucogranite intrusions in extensional regime of unroofing kinematics (VD2) in the time range of **355–345 Ma**. It is supposed that primary foliation was not so steep and their present spatial position was completed by Alpine (Cenozoic) AnD3 shearing. Dominant structural elements represent metamorphic foliations generally trending NE–SW (with variability in the NNE–SSW to ENE–WSW trend; Fig. 47) with rarely identified fold axes of the same direction and mineral lineations of metamorphic minerals with a shallow orientation on foliation surfaces (Fig. 48). From this, it is possible to infer the shear origin of the fold structures, or to the maximum component of the deformation stress, close to the direction of the fold axes. Based on the dating of monazites in paragneites and orthogneites, it can be




Fig. 47. Courses of metamorphic foliation in paragneisses (prevailingly the Suchý massif – 96 measurements): A – poles of planes, B – contourgram of the poles of the planes, C – prevailing trends of planes, D – dips of planes in paragneisses. Dominating foliations have NE–SW trend with steep dip to NW.

concluded that the maximum metamorphic overprint of rock complexes was around **380 Ma**, which corresponds to the Middle to Upper Devonian boundary and process over the subduction zone (**VD1s**).

Shear deformation in the ductile conditions of the amphibolite facies led, depending on the position of the rock complex in the continental crust, to the tectonic separation of leucogranite melt located in fold closures of migmatized paragneisses (Figs. 16 and 17), through the boudinage of leucogranite bodies situated in fold limbs still in supersolid to subsolid state up to the placement of thin leucogranite interbeds in semiductile flexures (Fig. 16). This indicates the decompression of the whole complex and gradual change of rheological conditions from ductile, to semiductile up to brittle conditions. Orogenic metamorphism, leading maximally to first manifestations of local melting of metamorphic complexes, it was followed by decompression and intrusion of thicker leucogranite volumes in shear regime. Granitoid melts were placed to higher crustal levels in ductile state, which is indicated by structures of magmatic flow, defined by position of dark minerals, but also feldspars. In more extreme cases podmienkach along the contact of leucogranite intrusions

with metamorphic mantle there occurred also synintrusive deformations of leucogranite, being manifested by origin of higher temperature foliation in conditions close to solidus, but also in subsolidus stage (this phenomenon was observed mainly NE of the Suchý vrch peak). These tectonites were distinguished also in geological map as syntectonic leucogranites with preferred orientation.

The structural analysis indicates that only part of inhomogeneities in granitoids can be derived from relic “pre-granitoid” setting of metamorphites and bigger part corresponds to own dynamics of magmatic flow of leucogranites, which is moderately counter-clockwise rotated with respect to older metamorphic foliation (Figs. 48 and 49).

Regional metamorphic structures (**VD1s**, **VD1c**) were overprinted by the heat of intruding granite masses (**VD2**, **MV2**). This overprint produced large macroscopically decipherable biotite II porphyroblasts (Bt of 2nd generation), though also these crystallized with preferred orientation due to acting stress field. Thermal effect of close leucogranite intrusions manifests in microscale by the origin of red-brown biotite and formation of numerous small garnets.




Fig. 48. Lineation of biotite in paragneisses (21 measurements) manifest the presence of two lineations. The more distinct of them trending ENE–WSW we relate to formation of bigger, better measurable biotites of 2nd generation, connected with thermal effect of penetrated leucogranite intrusions. Lineations trending NW–SW are probable connected with biotite of orogenic metamorphic genesis. Dominating shallower lineations indicate acting shear regime at penetrating leucogranite bodies.

Chemical dating of monazite revealed the age **355–345 Ma** of maximum intrusive activity and calculation of its PT conditions indicates pressure 3-4 kbar (for the stage of syntektonic S-type leucogranites **352–351 Ma**).

The shear emplacement of S-type leucogranite intrusions under decompression regime conditions (**VD2**) was followed with later opening of deeper crustal structures used by the intrusion of I-type omnidirectional biotitic granodiorites. They contain fine-grained mafic globules (currently not examined further; Fig. 41), corresponding with mafic portions of the magma, as well as rare rocks of (gabbro-)dioritic composition. This type of intrusion has a linear course (**VD4?**) and crops out oblique to older structures in NE part of Suchý massif. By this way the transition from S-type granite intrusions of exclusively crustal origin to more mafic intrusions of I-type reflects **the increase of thermality of granite melt towards the end of granite magmatism**. It indicates the deeper crustal, resp. subcrustal genesis of later magmatic melts. Such time succession, where the I-type granitoid intrusions in short time sequence followed after S-type granites

was demonstrated in granitoids of the Malé Karpaty Mts (Kohút et al., 2009).

In our work we used U-Th-Pb chemical dating of monazite, applying methodology by Konečný et al. (2018), which after statistic processing of its results can provide results close to zircon dating. It is valie especially at rapidly cooling magmatic systems without additional influence of the Mnz crystals by the fluid regime. With respect to proved Variscan decompression in the crystalline basement of the Suchý massif it is probable that Mnz dating provided data close to real intruive age of granites. It is valit mainly for leucogranites, connected with the phase of their rapid shear emplacement (**352–351 Ma**) and formation of pegmatite veins, finishing magmatic regime (**343–345 Ma**; Tournaisian / Visean boundary).

At I-type granodiorites, which finish magmatic process from geological and structural viewpoint, this age is probable influenced by the presence of older monazites, reflecting the mixed origin of the monazite crystals.

The group of S-type granitoids, indicating the contamination with surrounding paragneisses, the mixed




Fig. 49. Foliations (schliers – planar orientation of biotite, eventually feldspars) in granitoids (36 measurements): A – poles of planes, B – contourgram of the poles of planes, C – prevailing trends of planes, D – dips of inhomogeneities in granitoids. The contour diagram of planes poles (B) shows two distinct maxima, more often corresponding to planes trending NNE–SSW with steep dip to WNW, resp. ESE. Second, less distinct maximum corresponds to planes trending ENE–WSW with steep dip to SSE resp. NNW. While second, less distinct maximum probable represents relict structures connected with metamorphic rim of intrusion, first – very distinct maximum reflects the penetration dynamics of leucogranite magma.




Fig. 50. Prevailing courses of planar structures, filled with pegmatite and aplite-pegmatite melt (30 measurements), representing the last stage of leucogranite intrusions. They have prevailing N–S course with steep dip to W, less NE–SW and E–W. Relatively steep disjunctive structures formed in the final extension phase of the generating of granitoids. A – poles of planes, B – contourgram of the poles of the planes, C – prevailing trends of planes, D – dips of pegmatite and aplite veins.




Fig. 51. Diagram of the opening of space for leucogranite intrusions. A – the beginning of the shear regime with the deformation of the metamorphic complex; B – diagram of space opening in a more advanced stage of the shear deformation; C – block diagram of syn-deformation leucogranite intrusions (omnidirectional up to indistinct preferential orientation of minerals – red crosses; and syn-intrusively oriented marginal parts of intrusions – red crosses and dashes). Formation of thermally induced biotite II (brown ellipses). Thermally (granite thermal effects) slightly, or no affected complex of paragneiss. Transversely steep structures filled with aplites and zonal pegmatites, formed from leucogranite magma (orange). Black arrows show the sense of tectonic shear transport.

origin of monazites, reflects the age of intrusion as well as the age of formation of older monazites, analogical with those in paragneisses (which probable reflect the age of orogenic (regional) metamorphic process and in smaller scale also the age of protolith).

SHRIMP dating of zircon of S-type two-mica granite from Magura part of the Strážovské vrchy Mts (Kohút & Larionov, 2021) has shown two age modal groups of zircons (sample MM-29), older, with Neoproterozoic (Cadomian) age **568–551 Ma** and younger **365–361 Ma** (Devonian / Lower Mississippian) with Concordia age **360,9 ± 2,7 Ma**.

Older assumptions and measurements, indicating that I-types granitoids are significantly younger (**310–303 Ma**; Bibikova et al., 1990; Broska et al., 1990) than S-types, in the W. Carpathians were not confirmed by subsequent zircon dating. The LA-ICP-MS zircon dating from Kriváňska Malá Fatra (Broska & Svojtka, 2020) indicated only a small age difference among zircons from S-type granites (**342 ± 3 Ma**) and zircons from I-type granitoids (cores with magmatic age of **353 ± 3 Ma** and rims **342 ± 3 Ma** – this age was probably influenced by fluid regime?), so the I-type granitoids should be slightly older.

On the contrary, the SHRIMP zircon dating (Kohút et al., 2009) in S-type granites of the Bratislava massif (**355 ± 5 Ma**) and I-type tonalites of the Modra massif (**347 ± 4 Ma**) in Malé Karpaty Mts pointed out a small age difference between two large groups of granitoids of W. Carpathians, however, I-type granitoids appear younger. Just like in the Malé Karpaty Mts (Kohút et al., 2009), also in Strážovské vrchy Mts, it is confirmed that S-types and I-types of granitoids were produced in the same collision event, but they represent different stages of decompression – shearing regime (S-types) until the opening of the space in the final, extensional phase of the orogenic cycle, which was associated with the output of higher thermal, deeper-based magmas of I-type, with manifestations of hybridism, with accompanying diorite magmas of small volumes.

According to the chemical dating of the monazite in the Strážovské vrchy, the age of I-type granites intrusion is older than the age of the monazite of the S-type granites associated with the shear-deformation regime. Geological relationships, however, indicate the opposite. I-type intrusives penetrate mainly in the E-W direction (ENE–WSW; i.e. at a small angle) through older deformation structures, which are associated with the emplacement of S-type granites in the NE-SW direction in a shear regime. Thus, I-type granodiorites can be considered late- to post-tectonic in relation to shear deformation associated with syntectonic intrusions of leucogranites.

Therefore, we assume that the intrusions of I-type granodiorites-tonalites and rarely diorites should be close in age to the formation of pegmatites, or follow them (within the Tournaisian / Viséan interface). For this reason,

it is necessary to carry out a geochronological investigation of zircon in I-type granitoids from this area.

The interpretation of the late emplacement of I-type granodiorites and small diorite bodies in the Strážovské vrchy is in agreement with the statement by Kohút & Larionov (2021): “The peak at ca 342–340 Ma, representing probably the main period of the collisional granites production / emplacement, was in part accompanied by intrusions of dioritic syn-plutonic dykes.”

We determined the age of the change in deformation conditions in metamorphic complexes from ductile to semi-ductile and brittle conditions, based on monazite dating to approximately 345 Ma, which is the typical age of emplacement of aplite and pegmatite veins in brittle structures. Aplite and pegmatite veins are steeply dipping (mainly N–S dip) and their trend is generally oriented at a large angle to the main shear stress component (dominantly E–W direction), as was found in the eastern part of the Suchý massif. The mechanism of emplacement of granites in shear zones during the extensional regime has received more attention since the 1990s (Hutton et al., 1990). It is a mechanism when thickened continental crust is decompressed with the simultaneous ascent of granites in shear zones parallel to the axis of the orogen (orogen-parallel extension). Such placement mechanism is known e.g. in Caledonides (Braathen et al., 2000), Variscides (Oberc-Dziedic et al., 2015), or in “younger” ranges, e.g. from Greater Himalaya (Xu et al., 2013), or nearby Alpine Periadriatic line with several intrusive complexes (Márton et al., 2006). In the environment of the Variscan shear zone, a transition from metatexites to diatexites accompanied by an increase in the volume of granite mobilizes up to the placement of peraluminous granodiorite in the shear zone was demonstrated (De Luca et al., 2023).

Later intrusions of I-type granitoids are evidence of the increase in thermality of the granitoid process at the end of the decompression granitogenic stage. A suitable mechanism for such a process is a model of the subsidence of a relict block of oceanic crust after a collision into the asthenosphere (slab breakoff), thermal ascent of mantle masses, acceleration of decompression of crustal parts, its melting associated with the ascent of deeper-seated intrusions. (napr. von Blanckenburg & Davies, 1995). For granitoids of the Low Tatra Mts, such a model is discussed by Maraszewska et al. (2022).

Orthogneiss represent a special phenomenon within the Suchý crystalline basement. Less significantly they occur also in the western part of the Magura massif. From the viewpoint of Variscan setting we consider them as the upper structural element. On the older geological map of Mahel' (1982) they were considered as migmatites. The presence of spindle-shaped polycrystalline glomeroblasts composed of feldspars and quartz, which are a manifestation of ductile deformation of older phenocrysts

and the absence of leucosome formation, testify against such interpretation. We assume that this crustal element (which, due to the monotonous character, composition and dating of the metamorphic reworking, we consider to be a pre-Variscan granitoid; pre-VD0), within Variscan collision being obducted (VD1o) onto those parts of crystalline basement that primarily sedimented on the Paleozoic continental slope, or even on the ocean floor.

Orthogneiss lithologies were identified as a common component within the Western Carpathian crystalline basement. Several authors have investigated their age characteristics based on zircon dating. In the Western Tatras, they were dated from the upper and lower tectonic units using the method LA-MC-ICP-MS method by Burda & Klötzli (2011) to 534 Ma, which they considered to be the age of the protolith, while the age of 387 Ma they consider as the age of highly metamorphic Eo-Variscan event. The age of the protolith of the Veporic orthogneisses

from Muráň (Gaab et al., 2005) was determined to 464 ± 34 Ma (later redefined by Putiš et al., 2008 to Cadomian). Isochrone age of zircons from orthogneiss from the Low Tatra Mts (sample NTJ-1) was determined by Putiš et al. (2003) to 381 Ma, whereas model ages for different size fractions of zircons vary for isotopic system $^{207}\text{Pb}/^{235}\text{U} = 466.2\text{--}576.6$ Ma and for isotopic system $^{206}\text{Pb}/^{238}\text{U} = 441.2\text{--}526.4$ Ma. From this, it can also be concluded that the age of 381 Ma represents the high-temperature metamorphic overprint of the protolith. Later dating of orthogneisses, but also banded amphibolites from W. Carpathians (Putiš et al., 2008) provided wider Neo-Proterozoic-Cadomian age extent (640–530 Ma) at larger group of orthogneisses. Our chemical dating of monazite from orthogneisses with a distinct peak of 390 Ma can undoubtedly be considered as the metamorphic age of monazite, while ages in the range of 340–350 Ma represent the age of a subsequent thermal event associated with leucogranite intrusions.




Fig. 52. Planar elements in orthogneisses (28 measurements): A – poles of plains, B – contourogram of poles of plains, C – prevailing trend of planes, D – dips of metamorphic schistosity of orthogneisses.

The metamorphic schistosity of orthogneisses trending dominating ENE–WSW with steep to moderate dip to SSE–JJV. The field configuration indicates the shallow fan-like setting of orthogneiss body. The presence of distinct biotite lineations indicates a significant stress field acting at emplacement of orthogneisses in the Variscan setting. The more external (higher) position in the Variscan setting is emphasized by the absence, or the rarity of granitization manifestations, while transversal brittle structures in orthogneisses are filled only with aplitic and pegmatite veins (Figs. 24A and 39A).

When analyzing the structural elements of the crystalline basement in megascale, it is necessary to state that these very probably demonstrate Alpine-modified directions, which can be derived from the variable and often steep dips of the Mesozoic cover of Tatricum (Malá Magura cover succession) towards the NW. The Alpine tilting of the crystalline basement in the NW direction would indicate that the crystalline structures were significantly more gently dipping to the NW in the pre-Alpine period.

Regarding the configuration of the entire structure of the Magura and Suchý massifs, it is necessary to point out again that the western part of the Strážovské vrchy crystalline basement – the Suchý massif represents more external and less granited part of Variscan setting, manifested mainly by sharp intrusive contacts of granitoids to metamorphites. In the western part of the Suchý massif, approximately from the Radiša valley, surfaces of metamorphic foliation have shallower dip, being accompanied with diaphoresis of former metamorphic minerals from the greenschists facies. In microscale it is manifested by cataclasis of garnet phenocrysts, its new rotation, chloritization along cataclastic fractures, forming of actinolite at the expense of common amphibole in metabasites and replacement of original Al silicates in metapelites by glomeroblasts of fine-grained muscovite. We associate new metamorphic schistosity and retrograde mineral replacements with Late Variscan unroofing in shallow crustal levels.

Alpine metamorphic overprint

Alpine overprint (ApD, AnD) is not always reliably distinguishable from Late Variscan unroofing (VD2), however, there are some diagnostic signs that distinguish them. Fission-track dating of zircon from the Tatric crystalline basement (Marko et al., 2017) indicates that this basement during Alpine cycle was not overprinted by temperature higher than 320 °C (zircon closure temperature), which is valid also for sediments of the Malá Magura unit. Younger, Alpine low temperature metamorphism in the Magura massif (Čík & Petřík, 2014) was characterized by the presence of margarite and pumpellyite. Calculations of Alpine association in

the Poruba area in migmatite provided values 480 °C at pressure 4.6 kbar (which we consider an unrealistically high value) and in paragneiss 300 °C at pressure 2.9 kbar. Radial arranged pumpellyite crystals can occasionally be found in granite fractures also in Suchý massif and often in thin sections from metabasites, mainly in W margin of the Suchý massif.

Planes of Alpine cleavage in crystalline basement have prevailing trends ENE–WSW VSV–ZJZ (55–65°) with dominant dip to SE, less to NW. The linear kinematic indicators show mainly to horizontal shifts (strike-slips) related to the Paleo-Alpine thrusts of higher units (ApD1c), mainly of the Fatricum, which was partially reflected also in the Tatric basement.

Thermal modelling (Marko et al., 2017) based on fission-track dating of zircon (ZFT ages, reflecting thermal transition between 230–250–max. 320 °C) indicated ages corresponding to gradual ascent of Paleozoic consolidated crystalline block, while the absence of Late Paleozoic sediments in Tatricum reflects denudation of Tatric segment down to the level of crystalline basement in the lowest Triassic. Younger ZFT ages (245, 239 and 222 Ma) reflect gradual, but irregular cooling of crystalline blocks in the Middle to Upper Triassic, possibly repeated uneven submergence of the crystalline basement during the Lower to Middle Jurassic (ages 198 and 168 Ma).

The distinguishing criterion for Late Variscan unroofing (VD2) and Alpine metamorphism (MAp) is mainly Alpine mineral metamorphism with the formation of pumpellyite facies minerals in metabasites, indicating lower temperature metamorphic conditions in comparison with Late Variscan diaphoresis. From the spatial view, the Alpine metamorphism is often bound with marginal parts of the massif in contact with sediments of Mesozoic Malá Magura unit, which are also strongly tectonically reduced. This is manifested by narrow mylonite zones parallel with the margins of the massif, which, in the case of binding to sediments with black schists (e.g. in the Čavoj area), are a source of metals remobilized into veins or stockworks in these narrow zones. During later Alpine updoming of crystalline basement after the completion of formation of the nappe setting, probably in the Upper Cretaceous or Paleogene, locally started in upper crustal conditions the origin of zones of low-temperature ultracataclasis (kakiritization), demonstrated by origin of dark afanitic zones thick up to several cm. At a later stage, during the continuing structuring of the crystalline blocks, wider zones of ultracataclasis were formed in the granitoids and quartzites, which are mainly present in the area of the Radiša fault (more appropriate name – Radiša fault zone), Závadka fault, Štovo fault, possibly also Diviaky fault (more appropriate name – Diviaky fault zone; Fig. 1). The final stage is the disintegration of the crystalline massif along the E-W oriented faults with the subsidence of the

southern blocks, as well as faults and fault zones in the N–S direction (e.g. Diviaky fault and Radiša fault; AnD4).

Acknowledgment

The authors appreciate the contribution of the research on the project *Geological map of the Strážovské vrchy Mts – Easten part* funded by the Ministry of Environment of the Slovak Republic. Results of petrological research by Rastislav Demko (State Geological Institute of Dionýz Štúr – ŠGÚDŠ Bratislava, Slovakia) are greatly appreciated. This paper is also a contribution of the State Geological Institute of Dionýz Štúr – ŠGÚDŠ, for the EC – CINEA HORIZON-CL5-2021-D3-D2 project 101075609 Geological Service for Europe (GSEU) within WP6 – Geological framework for the European geological data & information system and task T6.2.4 Lithotectonic units.

References

- BIBIKOVA, E. V., KORIKOVSKY, S. P., PUTIŠ, M., BROSKA, I., GOLZMAN, Y. V. & ARAKELIANTS, M. M., 1990: U/Pb, Rb/Sr and K/Ar dating of Sihla tonalites of Vepor Pluton (Western Carpathians). *Geologický zborník, Geologica Carpathica*, 41, 427–436.
- BRAATHEN, A., NORDGULEN, Ø., OSMUNDSEN, P. T., ANDERSEN, T. B., SOLLI, A. & ROBERTS, D., 2000: Devonian, orogen-parallel, oposed extension in Central Norwegian Caledonides. *Geology*, 28, 7, 615–618.
- BROSKA, I., BIBIKOVA, E. V., GRACHEVA, T. V., MAKAROV, V. A. & CAÑO, F., 1990: Zircon from granitoids rocks of the Tribeč-Zobor crystal-line complex: its typology, chemical and isotopic composition. *Geologický zborník, Geologica Carpathica*, 41, 393–406.
- BROSKA, I. & SVOJTKA, M., 2020: Early Carboniferous successive I/S granite magmatism recorded in the Malá Fatra Mountains by LA-ICP-MS zircon dating (Western Carpathians). *Geologica Carpathica*, 71, 5, 391–401.
- BURDA, J. & KLÖTZLI, U., 2011: Pre-Variscan evolution of the Western Tatra Mountains: new insight from U-Pb zircon dating. *Mineral. Petrol.*, 102, 99–115.
- DE LUCA, M., RUBERTI, N., OGGIANO, G., ROSSI, PH., PASCUCCI, V. & CASINI, L., 2023: Structure of a Variscan migmatite-granite transition zone (N Sardinia, Italy). *Journal of Maps*, 19, 1, 1–10. DOI:10.1080/17445647.2023.2182721.
- DYDA, M., 1988: Retrograde processes in paragneisses of the Suchy and Mala Magura Mts. *Geologický zborník, Geologica Carpathica*, 39, 241–257.
- DYDA, M., 1990: Metamorphic processes in paragneisses from the Suchy and Mala Magura Mts. (The West Carpathians). *Geologický zborník, Geologica Carpathica*, 41, 315–334.
- GAAB, A. S., JANÁK, M., POLLER, U. & TODT, W., 2006: Alpine reworking of Ordovician protoliths in the Western Carpathians: Geochronological and geochemical data on the Muráň Gneiss Complex, Slovakia. *Lithos*, 87, 261–275.
- HOVORKA, D. & FEJDI, P., 1983: Garnets of peraluminous granites of the Suchý and the Malá Magura Mts. (The Western Carpathians) – their origin and geological significance. *Geologický zborník, Geologica Carpathica*, 34, 1, 103–115.
- HOVORKA, D. & MÉRES, Š., 1991: Pre-upper carboniferous gneisses of the Strážovské vrchy upland and the Malá Fatra Mts. (The Western Carpathians). *Acta geogr. geol. Univ. Comen.*, 46, 103–169.
- HRAŠKO, L., PELECH, O., SENTPETERY, M., OLŠAVSKÝ, M., KRONOME, B., LAURINC, D., MAGLAY, J., NÉMETH, Z., BOOROVÁ, D., ŽECOVÁ, K., ZLINSKÁ, A., MIKUDÍKOVÁ, M., FEKETE, K., NAGY, A., DEMKO, R. & KOVÁČIK, M., 2020: Geologická mapa regiónu Strážovské vrchy – východná časť v mierke 1 : 25 000, listy 35-22-1 Pružina, 35-22-3 Valaská Belá, 35-24-1 Nitrianske Rudno a 35-24-3 Uhrovec. *Manuscript. Bratislava, archive ŠGÚDŠ*, 289 p.
- HRAŠKO, L., KOVÁČIK, M. (eds.), OLŠAVSKÝ, M., SENTPETERY, M., PELECH, O., LAURINC, D., MAGLAY, J., NÉMETH, Z., KRONOME, B., NAGY, A., KOVÁČIK, M.†, VLAČIKY, M. & DANANAJ, I., 2021: Geologická mapa Strážovských vrchov (východná časť) 1 : 50 000, Bratislava, MŽP SR – ŠGÚDŠ. https://www.geology.sk/wp-content/uploads/documents/foto/geol_mapy_50k/58_StrazovskeVrchy_final.jpg.
- HUTTON, D. H. W., DEMPSTER, T. J., BROWN, P. E. & BECKER, S. D., 1990: A new mechanism of granite emplacement: intrusion in active extensional shear zones. *Nature*, 343, 452–455.
- IVAN, P. & MÉRES, Š., 2015: Geochemistry of amphibolites and related graphitic gneisses from the Suchý and Malá Magura Mountains (central Western Carpathians) – evidence for relics of the Variscan ophiolite complex. *Geologica Carpathica*, 66, 5, 347–360.
- IVANOV, M., 1957: Genéza a vzťah granitoidných intrúzií k superkrustálnym sériám kryštalinika Suchého a Malej Magury. *Geologické práce, Zošit*, 47, 87–115.
- KAHAN, Š., 1976: Štruktúrna charakteristika kryštalinika masívu Suchého a Malej Magury. *Manuscript. Bratislava, archive Katedry geológie a paleontológie PFUK*.
- KAHAN, Š., 1979: Čiastková správa o výskume kryštalinika Strážovských vrchov, riešenie v rokoch 1976–1978. *Manuscript. Bratislava, archive ŠGÚDŠ (arch. no. 45428)*, 81 p.
- KAHAN, Š., 1980: Strukturelle und metamorphe Charakteristik des Gebirges Strážovské vrchy (Suchý und Malá Magura). *Geologický zborník, Geologica Carpathica*, 31, 4, 577–602.
- KAHAN, Š., ZELMAN, J. & PUTIŠ, M., 1978: Správa o prácach vykonaných v rámci HZ 23/74 – VČ a HZ 36/76 VČ (Suchý – Malá Magura) v r. 1974–1976. *Manuscript. Bratislava, archive ŠGÚDŠ (arch. no. 50327)*, 134 p.
- KLINEC, A., 1956: Zpráva o mapovaní kryštalinika pohoria Malá Magura. *Manuscript. Bratislava, archive ŠGÚDŠ*.
- KLINEC, A., 1958: Kryštalinikum severovýchodnej časti Malej Magury. *Geologické práce, Správy*, 12, 93–101.
- KOHÚT, M., UHER, P., PUTIŠ, M., ONDREJKA, M., SERGEEV, S., LARIONOV, A. & PADERIN, I., 2009: SHRIMP U-Th-Pb zircon dating of the granitoid massifs in the Malé Karpaty Mountains (Western Carpathians): evidence of Meso-Hercynian successive S- to I-type granitic magmatism. *Geologica Carpathica*, 60, 345–350.
- KOHÚT, M. & LARIONOV, A. N., 2021: From subduction to collision: Genesis of the Variscan granitic rocks from the Tatric Superunit (Western Carpatians, Slovakia). *Geologica Carpathica*, 72, 2, 96–113.

- KONEČNÝ, P., KUSIAK, M. A. & DUNKLEY, D. J., 2018: Improving U,Th,Pb electron microprobe dating using monazite age references. Invited review article. *Chemical Geology*, 484, 22–35.
- KRÁL, J., GOLTZMAN, Y. & PETRÍK, I., 1987: Rb-Sr whole-rock isochron data of granitic rocks from the Strážovské vrchy Mts: The preliminary report. *Geologický zborník, Geologica Carpathica*, 38, 2, 171–179.
- KRÁL, J., HESS, C. KOBER, B. & LIPPOLT, H. J., 1997: $^{207}\text{Pb}/^{206}\text{Pb}$ and $^{40}\text{Ar}/^{39}\text{Ar}$ age data from plutonic rocks of the Strážovské vrchy Mts. basement, Western Carpathians. In: Grecula, P., Hovorka, D. a Putiš, M. (eds.): Geological evolution of the Western Carpathians. *Mineralia Slovaca, Monogr.*, 253–260.
- MAHEE, M. & GROSS, P., 1975: Vysvetlivky mapy 1 : 25 000 listy Kšinná, Nitrianske Rudno (sčasti i Nitrianske Sučany) – južná časť Strážovskej hornatiny, čiastková záverečná správa za rok: 1975. *Manuscript. Bratislava, archive ŠGÚDŠ*, 1–39.
- MAHEE, M., 1983: Vysvetlivky ku geologickej mape Strážovských vrchov v mierke 1 : 50 000. *Bratislava, Geologický ústav Dionýza Štúra*, 90 p.
- MAHEE, M., 1985: Geologická stavba Strážovských vrchov. *Bratislava, Geologický ústav Dionýza Štúra*, 221 p.
- MARASZEWSKA, M., BROSKA, I., KOHÚT, M., YI, K., KONEČNÝ, P. & KURYLO, S., 2022: The Ďumbier-Prašivá high K-calc-alkaline granite suite (Low Tatra Mts., Western Carpathians): Insights into their evolution from geochemistry and geochronology. *Geologica Carpathica*, 73, 4, 273–291.
- MARKO, F., ANDRIESEN P. A. M., TOMEK, Č., BEZÁK, V., FOJTÍKOVÁ, L., BOŠANSKÝ, M., PIOVARČI, M. & REICHWALDER, P., 2017: Carpathian Shear Corridor – a strike-slip boundary of an extruded crustal segment. *Tectonophysics*, 703–704, 119–134.
- MÁRTON, E., TROJANOVA, M., ZUPANIČIČ, N. & JELEN, B., 2006: Formation, uplift and tectonic integration of a Periadriatic intrusive complex (Pohorje, Slovenia) as reflected in magnetic parameters and palaeomagnetic directions. *Geophys. Jour. Internat.*, 167, 3, 1148–1159.
- MIKOLÁŠ, S., KOMORA, J., KANDERA, K., SANDANUS, M., ŠLEPECKÝ, T., JANUŠ, J., STAŇA, Š., OČENÁŠ, D. & STUPÁK, J., 1995: Čavoj – Gápel – Ag, Pb, Zn rudy – vyhladávací prieskum, stav k 31. 3. 1995. *Manuscript. Bratislava, archive ŠGÚDŠ (arch. no. 82971)*, 114 p., 44 príl.
- MONTEL, J. M., FORET, S., VESCHAMBRE, M., NICOLLET, CH. & PROVOST, A., 1996: Electron microprobe dating of monazite. *Chemical Geology*, 131, 1–4, 37–53.
- NÉMETH, Z., PUTIŠ, M. & HRAŠKO, L., 2016: The relation of metallogeny to geodynamic processes – the natural prerequisite for the origin of mineral deposits of public importance (MDoPOI: The case study in the Western Carpathians, Slovakia). *Mineralia Slovaca*, 48, 2, 129–135.
- NÉMETH, Z., 2021: Lithotectonic units of the Western Carpathians: Suggestion of simple methodology for lithotectonic units defining, applicable for orogenic belts world-wide. *Mineralia Slovaca*, 53, 81–90.
- OBERC-DZIEDZIC, T., KRYZA, R. & PIN, CH., 2015: Variscan granitoids related to shear zones and faults: examples from Central Sudetes (Bohemian Massif) and the Middle Odra Fault Zone. *Int. Jour. of Earth Sci.*, 104, 1139–1166.
- PUTIŠ, M., KOTOV, A. B., PETRÍK, I., MADARÁS, J., SALNIKOVA, E. B., YAKOVLEVA, S. Z., BEREZHNYAYA, N. G., PLOTKINA, Y. V. KOVACH, V. P., LUPTÁK, B. & MAJDÁN, M., 2003: Early- vs. Late orogenic granitoids relationships in the variscan basement of the Western Carpathians. *Geologica Carpathica*, 54, 3, 163–174.
- PUTIŠ, M., SERGEEV, S., ONDREJKA, M., LARIONOV, A., SIMAN, P., SPIŠIAK, J., UHER, P. & PADERIN, I., 2008: Cambrian-Ordovician metaigneous rocks associated with Cadomian fragments in the West-Carpathian basement dated by SHRIMP on zircons: a record from the Gondwana active margin setting. *Geologica Carpathica*, 59, 1, 3–18.
- VIELZEUF, D. & MONTEL, J. M., 1994: Partial melting of metagreywackes: fluid-absent experiments and phase relationships. *Contrib. Mineral. Petrol.*, 117, 375–393.
- VILINOVIČOVÁ, L., 1988: Geochémia a petrogenéza granitoidných hornín Suchého a Malej Magury a ich vzťah k rulám. Kandidátska dizertačná práca. *Manuscript. Bratislava, archive ŠGÚDŠ (arch. no. 68031)*, 64 p.
- VILINOVIČOVÁ, L., 1990: Petrogenesis of gneisses and granitoids from the Strážovské vrchy Mts. *Geologický zborník, Geologica Carpathica*, 41, 4, 335–376.
- WHITE, R. W., POWEL, R. & JOHNSON, T. E., 2014: The effect of Mn on mineral stability in metapelites revised: New a-x relations for manganese-bearing minerals. *Jour. Metamorphic. Geol.*, 32, 8, 809–828.
- WU, CHUN-MING, ZHANG, J. & LIU-DONG REN, 2004: Empirical Garnet-Biotite-Plagioclase-Quartz (GBPQ) geobarometry in Medium- to High-Grade Metapelites. ??????, 45, 1907–1921.
- XU, ZH., WANG, Q., PÉCHER, A., LIANG, F., QI, X., CAI, ZH., LI, H., ZENG, L. & CAO, H., 2013: Orogen-parallel ductile extension and extrusion of the Greater Himalaya in the late Oligocene and Miocene. *Tectonics*, 32, 191–215.
- VON BLANCKENBURG, F. & DAVIES, J. H., 1995: Slab breakoff: A model for syncollisional magmatism and tectonics in the Alps. *Tectonics*, 14, 1, 120–131.

Variské litotektonické jednotky v masíve Suchého v Strážovských vrchoch, Západné Karpaty – výsledok sedimentárnych, tektonometamorfných a granitizačných procesov

V predalpínskej štruktúre západnej časti Strážovských vrchov (masív Suchého) boli vyčlenené tri litologicky odlišné variské litotektonické jednotky: 1. sedimenty, ktoré majú pôvod v hlbšom oceánskom bazéne (prevažne metapelity s rôznym obsahom organickej hmoty, metabazalty, metakarbonáty?); 2. sedimenty kontinentálneho svahu (flyšoidné sedimenty s prevahou drobových sedimentov); 3. jednotka kontinentálneho pôvodu blízka granitovému zloženiu (ortoruly), ktorá má pravdepodobne predvariský pôvod. Tieto komplexy rôznej geotektonickej proveniencie boli amalgamované a metamorfované pred granitizačným štádiom (štádium pre-VmD2 pred obdobím misisipu). Z pohľadu variskej polyorogénnej evolúcie variské procesy v tatriku reprezentujú mezovariskú evolúciu (VmD).

Maximálne P-T podmienky orogénnej (regionálnej) metamorfózy (do 610 °C a 7,5 – 8,5 kbar) neboli dostatočné na rozsiahlejšie prejavy anatexie. Z terénnych pozorovaní vyplýva, že k produkcii obmedzeného objemu granitoidných tavenín dochádzalo najmä pri kontaktoch amalgamovaných litotektonických jednotiek, pravdepodobne s prispením strižného tepla.

Pravdepodobne vplyvom dodatočného tepla produkovaného pod akrečným klinom spolu s dekompresným režimom nastala etapa tvorby a umiestňovania granitoidov v orogénnej fáze VmD2.

Etapa tvorby granitov počas misisipu (štádium VmD2) je spojená s umiestňovaním rôznych typov granitoidných magiem, pričom ako geologicky najstaršie vystupujú granodiority s častými šlírmami. Predstavujú málo diferencovanú a málo mobilnú kryštalovú kašu (jv. časť územia).

V hlavnom deformačnom štádiu súhlasne s deformačným plánom v metamorfovaných komplexoch intrudovali masy leukogranitov, ktoré interagovali s okolitými metamorfovanými horninami v plytších kôrových podmienkach a spôsobili kontaktnú premenu staršej metamorfnej asociácie (do 590 °C a 3 – 4 kbar). Syndeformičný charakter leukokratných granitov je daný usmernením lupenňov biotitu paralelne s deformačným plánom v okolitých metamorfitoch. Časť leukogranitov najmä v centrálnych častiach telies je všesmerná. Deformačné pole pôsobilo

nielen počas intrúzie leukogranitovej magmy, ale aj v sub-solidovom štádiu.

Záverečnou etapou tohto procesu bola tvorba veľkých telies pegmatitov v extenzných fraktúrach štádia VmD2, orientovaných pod veľkým uhlom k hlavnej zložke napätia. Textúra pegmatitov so sivým blokovým K-živcom a nedostatok minerálov obsahujúcich vodu poukazuje na pravdepodobnú pneumatolytickú frakturáciu v následne otvorenom prostredí štádia VmD2.

Záverečnou etapou štádia VmD2 sú intrúzie granodioritov typu I so znakmi miešania magiem (*mixing*, resp. *mingling*).

Chemické datovanie monazitu v granitoidoch umožňuje datovať jednotlivé fázy granitizačného procesu medzi 360 – 345 mil. rokov, pričom najmladšie veky korešponujú so vznikom pegmatitov. Datovanie monazitu v metamorfovaných horninách poukazuje na termálne ovplyvnenie granitizačným procesom (360 – 350 mil. r.) a zároveň indikuje staršie regionálne metamorfné prepracovanie prevažne v období 370 – 380 mil. rokov. Staršie reliktné údaje, ktoré je problematické interpretovať, môžu byť sčasti odvodené od veku protolitu metamorfovaných hornín.

Scenár umiestňovania granitoidných intrúzií je súhlasný s dekompresným režimom (v štádiu VmD2) po závere kôrového zhrubnutia (v štádiu VmD1c) až po frakturáciu kôrového bloku s intrúziami magiem typu I hlbšieho pôvodu.

Po tomto období pokračovala exhumácia blokov kryštalínika, čiastočná diaforéza a neskôr povrchová erózia až do spodného triasu.

Opätovné ponorenie kryštalinických komplexov je spojené s nízkym stupňom alpínskeho metamorfného prepracovania.

Doručené / Received: 30. 5. 2024
Prijaté na publikovanie / Accepted: 28. 6. 2024

Appendix

Table of sample basic field characteristic and coordinates

Sample	Field characteristics	WGS-84 coordinates			SJTSK coordinates		
		φ	λ		X	Y	
LHS-42	Contact of Bt paragneiss and pegmatite with grey blocky Kfs. Metamorphic schistosity 320/85.	48.820 482	18.416 903		1 214 869.74	470 428.22	
LHS-57	Metamorphically differentiated Bt paragneiss with small interbeds of granitic material, subvertical metamorphic schistosity in N-S direction.	48.816 606	18.441 510		1 215 450.29	468 663.70	
LHS-73	Stock of the Ms-Bt massive granite.	48.838 813	18.433 411		1 212 940.00	469 050.00	
LHS-94A	Medium-grained schlier granitoid with moderate preferred orientation of Bt+Plg, Kfs, contains planes rich in Bt.	48.839 33	18.448 381		1 212 974.39	467 950.35	
LHS-94B	V LHS-94A small body of Plg-rich (up to 50 %) granodiorite with abundant Bt. Idiomorphic Plg large up to 1–2 mm. Interstices are filled with Bt.	48.839 33	18.448 381		1 212 974.39	467 950.35	
LHS-114	Alternation of migmatites with predominant metamorphic schists with the direction of dip 300/85°. 2 generations of granites are present in them: 1. granite – ductile folded, parallel to the metamorphic schist. 2. In the fold cores there are coarse-grained Ms-Bt granites, with idiomorphic feldspars. This 2nd type is hybridized at the edges, finer-grained. The presence of Bt schists indicates low magma mobility. This 2nd type probably segregates into large volumes, so that granitoids predominate in the western flank of the outcrops.	48.837 475	18.464 977		1 213 281.35	466 753.53	
LHS-131	Ductile folded migmatites with a ratio of melt to metatect of 1 : 1. In this, concordant positions of medium-grained leucogranites with inhomogeneously distributed Bt are placed, in them the Bt is parallel to that in the migmatites.	48.837 114	18.465 415		1 213 323.96	466 724.77	
LHS-137	Stromatitic migmatite with 1 cm thick parallel granite positions.	48.831 848	18.467 765		1 213 921.89	466 601.54	
LHS-153	Fine-grained amphibolitic rocks, containing positions of leucogranite veins parallel to the metamorphic foliation, a few cm thick, up to a younger generation of transversal granite veins, which branch intricately. Veins of pegmatitic granites are also present.	48.840 364	18.462 299		1 212 944.81	466 922.69	
LHS-177-2	177-1 light-grey Bt paragneisses without foliation planes with Bt2, but Bt is lineated; 177-2 hybridic, 177-2 hybridic medium- to coarse-grained granodiorites with irregularly spatially and dimensionally arranged Bt (together with 177-1 light-grey Bt paragneisses without foliation planes with Bt2, Bt is lineated).	48.819 231	18.456 915		1 215 253.67	467 512.06	
LHS-185	Schlier granitoids with xenoliths of fine-grained Bt paragneiss.	48.836 248	18.459 216		1 213 382.14	467 186.30	
LHS-197	Grey, fine-grained paragneiss, schistose, which are affected by new thermal reworking, with the formation of larger Bt2 flakes. Metamorphic schistosity is subvertical in E-W direction.	48.836 381	18.397 549		1 212 988.88	471 695.77	
LHS-200	Tectonized, partially diaphtorized? medium-grained Bt + Ms leucogranites without paragneisses.	48.842 981	18.400 166		1 212 273.72	471 442.73	
LHS-204	Grey, metamorphically foliated fine-grained Bt paragneisses, with new thermal reworking, with 1–2 mm Bt2 flakes, parallel with new foliation surfaces.	48.828 532	18.389 992		1 213 811.65	472 326.31	

Tab. – continue

Sample	Field characteristics	WGS-84 coordinates		SJTSK coordinates	
		φ	λ	X	Y
LHS-206	Medium- to coarse-grained pegmatite leucogranite with transition to pegmatite. Several hundred m thick vein, cropping out in N side of the valley, where large blocks fall down to the stream. Locally present tiny Grt and grey Kfs. Grt forms subparallel interbeds, which proves the zonality. Ms in larger flakes usually forms nests.	48.828 998	18.432 599	1 214 022.56	469 200.63
LHS-207	Banded gneisses with metamorphic differentiation with straight flat alternating of 1 mm thick light and dark bands. Foliation planes are covered with larger Bt flakes. Transversally to metamorphic foliation penetrating fine- to medium-grained veinlets of leucogranite.	48.826 465	18.402 749	1 214 119.71	471 407.99
LHS-255	Paragneiss to micaschist with the formation of chlorite, in contact with aplite with garnet, metamorphic shale in the direction of dip 320/15°.	48.829 517	18.378 054 5	1 213 629.24	473 186.11
LHS-261	Grey aphanitic omnidirectional rock originally supposed to be metaquartzite. Contains Hbl.	48.846 793	18.378 087 0	1 211 715.29	473 021.80
LHS-272	Porphyroblastic orthogneiss with 5 mm porphyroblasts of rotated feldspars.	48.849 538	18.385 927 0	1 211 459.60	472 422.79
LHS-273	Porphyroblastic orthogneiss.	48.853 505	18.387 930 0	1 211 032.43	472 239.20
LHS-329	Dark and light grey, fine-grained banded paragneiss.	48.858 451	18.342 786	1 210 204.72	475 493.36
LHS-350	Dark grey Bt paragneiss with graphitic admixture, associated with black graphitic schists, schistosity dip 50/35°.	48.860 370	18.344 089	1 210 000.00	475 380.00
LHS-356a	Micaceous Bt paragneiss, metamorphic foliated structure, micas approx. 1 mm large (metapelite), contact with aplite.	48.837 683	18.367 388	1 212 658.42	473 889.70
LHS-356b	Micaceous Bt paragneiss, metamorphic foliated structure, micas approx. 1 mm large (metapelite), contact with aplite.	48.837 683	18.367 388	1 212 658.42	473 889.70
LHS-415	Oriented Bt-rich orthogneiss on the road base, metamorphic foliation, oblique pegmatite vein with grey Kfs.	48.863 594	18.423 373	1 210 132.55	469 553.65
LHS-482A	Bt grey paragneisses with steep metamorphic schistosity. In foliation planes occur large Bt ₂ , system is penetrated in shear regime by granites – leucocratic fine-grained with rare oriented Bt in magmatic flow, the same system is followed with coarse-grained to blocky pegmatites with grey blocky feldspars and rare Ms.	48.865 501	18.456 219	1 210 122.58	467 134.79
LHS-482B	Bt paragneisses with metamorphic schistosity, Bt ₂ , interbeds of leucogranite with inhomogeneously distributed Bt. The weakly bordered pegmatite is present in them, being zonal, thick to 10 m, in the middle with grey Kfs and Ab at the margin.	48.865 812	18.455 568	1 210 084.14	467 179.50
LHS-557	Two-mica granite with prevailing Ms and pegmatite.	48.837647	18.421309	1 212 995.11	469 946.15
LHS-572	In environment of Bt paragneisses with Bt ₂ , the grey to light-grey pegmatites frequently occur with sugar-type aplitic zone and often tiny Grt, central zone with Ms, Bt, Gtr, Kfs, Ab, Qtz.	48.849 571	18.407 010	1 211 585.96	470 880.79
LHS-599	Large blocks of pegmatites with blocky grey Kfs, applites with garnet. Rock contains only Bt, Ms is missing. Vein crops out in the environment of two-mica granites.	48.830 616	18.434 235	1 213 853.53	469 066.00

Tab. – continue

Sample	Field characteristics	WGS-84 coordinates		SJTSK coordinates	
		φ	λ	X	Y
LHS-607	Two-mica leucogranite with irregularly distributed, oriented Bt, and omnidirectional Ms, present are not sharply bordered pegmatites.	48.842 982	18.423 083	1 212 414.87	469 766.75
LHS-611	Large blocks of pegmatites with grey blocky K-feldspar in environment of leucogranites with oriented Bt.	48.827 868	18.424 823	1 214 100.00	469 780.00
LHS-623	Light deformed granites in subsolidic regime, planar orientation in outcrop, Bt with preferred orientation.	48.849 214	18.441 503	1 211 837.30	468 361.74
LHS-626	Light leucogranite with oriented Bt and feldspars, Kfs is large up to 2–3 mm.	48.849 991	18.434 700	1 211 709.53	468 852.01
LHS-650	Migmatite with a predominance of felsic component.	48.847 407	18.476 226	1 212 249.59	465 839.11
LHS-711	Orthogneiss with oriented biotite.	48.862 027	18.370 590	1 209 981.00	473 427.10
SZN-24	Light fine-grained rock, omnidirectional keratophyre?	48.841 015	18.368 883	1 212 298.32	473 749.04
SZN-25	Black metamorphosed schist, folded.	48.840 181	18.381 699	1 212 469.97	472 819.50
SZN-32a	Alternation of grey to black metamorphosed shales and metasediments.	48.844 398	18.365 499	1 211 902.50	473 964.71
SZN-34a	Dark metapelitic paragneiss with biotite, staurolite and garnet.	48.842 115	18.369 983	1 212 183.26	473 658.27
SZN-34b	Dark metapelitic paragneiss with biotite, staurolite and garnet.	48.842 115	18.369 983	1 212 183.26	473 658.27
SZN-34c	Dark metapelitic paragneiss with biotite, staurolite and garnet.	48.842 115	18.369 983	1 212 183.26	473 658.27
SZN-57	Dark-grey paragneiss with graphitic pigment.	48.861 108	18.354 930	1 209 985.55	474 580.53
SZN-66	Dark-grey to black paragneiss.	48.864 038	18.369 480	1 209 751.09	473 489.36
SZN-85	Fine-grained amphibolite.	48.845 178	18.374 989	1 211 874.84	473 263.40

Used abbreviations:

Minerals: Pl – plagioclase (Ab – albite, An – anorthite), Kfs – K-feldspar, Hbl, Amf – hornblende, Cpx – clinopyroxene, Grt – garnet (Alm – almandine, Spess – spessartine, Pyr – pyrope, Gross – grossular), Qtz – quartz, Bt – biotite, Ms – muscovite, Sill – sillimanite, Stau, St – staurolite, Chl – chlorite, Ep – epidote, Ap – apatite, Zrn – zircon, Mnz – monazite, Mag – magnetite, Ilm – ilmenite

Mg # – ionic ratio $Mg/(Fe_{total} + Mg)$

BSEI – back scattered electron images

TIMS – Thermal Ionization Mass Spectrometry

SHRIMP – Sensitive High Resolution Ion Micro-Probe

LA-ICP-MS – Laser Ablation Inductively Coupled Plasma Mass Spectrometry

Lithotectonic units – orogenic cycles, orogenic phases: VD – Variscan orogenic cycle, VmD0 – Meso-Variscan divergent orogenic phase, VmD1c – Meso-Variscan collisional phase, VmD2 – Meso-Variscan post-collisional phase with thermal overheating over hot line, extensional and unroofing kinematics, AD – Alpine orogenic cycle, AnD3 – Neo-Alpine orogenic phase characteristic with subhorizontal shearing kinematics, AnD4 – Neo-Alpine orogenic phase characteristic with regional extension and origin of pure shear-type faults preferably of E–W and N–E course.